



(11) EP 0 845 529 A2

EUROPEAN PATENT APPLICATION

- (12)  
(43) Date of publication:  
03.06.1998 Bulletin 1998/23  
(21) Application number: 97308562.4  
(22) Date of filing: 27.10.1997  
(51) Int. Cl. C: C12N 15/12, C07K 14/705, A61K 38/17, C07K 16/28, G01N 33/68

- (84) Designated Contracting States:  
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE  
(30) Priority: 29.10.1996 JP 288823/95  
(71) Applicant: TAKEDA CHEMICAL INDUSTRIES, LTD.  
Chuo-ku, Osaka 541 (JP)  
(72) Inventors:  
• Hinuma, Shuji, Takeda Chemical Industries Ltd.  
Chuo-ku, Osaka 541 (JP)  
• Fukusumi, Shoji, Takeda Chemical Industries Ltd.  
Chuo-ku, Osaka 541 (JP)  
(74) Representative: Connell, Anthony Christopher  
SmithKline Beecham plc  
Corporate Intellectual Property,  
Two New Horizons Court  
Brentford, Middlesex TW8 9EP (GB)  
Remarks:  
The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) Human G-protein coupled receptor protein cloned from fetal brain cDNA library

- (57) A novel G-protein coupled receptor protein, a partial peptide and their salts are disclosed. DNA encoding the receptor protein, production of the receptor protein, determination of a ligand to the receptor protein, a method for screening for compounds which inhibit lig-

Description

FIELD OF THE INVENTION

5 The present invention relates to novel G-protein coupled receptor proteins derived from a human brain, or salts thereof.

BACKGROUND OF THE INVENTION

10 Many hormones and neurotransmitters regulate functions in a living body through specific receptor proteins existing in a cell membrane. Many of these receptor proteins mediate signal transmission in a cell by activation of coupled guanine nucleotide-binding proteins (heterotrimer, sometimes, referred to as G-proteins) and are generally called G-protein coupled receptor proteins or 7-transmembrane receptor proteins because they contain a common structure having seven transmembrane domains.

15 G-protein coupled receptor proteins exist on each functional cell surface of cells and internal organs of a living body and play very important roles as targets of molecules which regulate functions of the cells and internal organs of the living body, for example, hormones, neurotransmitters, physiologically active substances and the like.

20 To clarify the relation between substances which regulate elaborate functions in cells and internal organs of various living bodies and their specific receptor proteins, in particular, G-protein coupled receptor proteins provide a very important means for clarification of functional mechanisms of cells and internal organs of various living body as well as for development of drugs having close relation to such functional mechanisms. For example, in a central nervous system organ such as a brain, its physiological functions are controlled through regulation by many hormones, hormone-like substances, neurotransmitters, physiologically active substances or the like. In particular, neurotransmitters are found in numerous sites within a brain and regulate the physiological functions through their corresponding receptor proteins.

25 However, it is supposed that many unknown neurotransmitters still exist in a brain and, as for their receptor proteins, many structures of cDNAs encoding such proteins have not yet been reported. In addition, it is still unknown if there are subtypes of known receptor proteins.

30 Also, to clarify the relation between substances which regulate elaborate functions in a brain and their specific receptor proteins provide a very important means for development of drugs. Further, for screening for agonists and antagonists to receptor proteins, efficiently, in development of drugs, it is required to clarify functional mechanisms of receptor protein genes expressed in a brain and to express them in a suitable expression system.

35 Recently, as a means for analyzing genes expressed in a living body, random analysis of cDNA sequences has been studied actively. The sequences of cDNA fragments thus obtained have been registered with data bases as Expressed Sequence Tags (ESTs) and are publicly available. However, for many of ESTs, it is difficult to deduce their functions from their sequential information only. For example, although two ESTs, accession No. T080399 (SEQ ID NO: 5) and No. T27053 (SEQ ID NO: 6), have been registered with the data base, NCBI dbEST, their functions are not clarified.

OBJECTS OF THE INVENTION

40 One object of the present invention is to provide novel G-protein coupled receptor proteins derived from a human brain, their partial peptides and their salts.

45 Another object of the present invention is to provide isolated DNA comprising DNA encoding the G-protein coupled receptor proteins or their partial peptides.

A further object of the present invention is to provide a recombinant vector comprising the DNA and a transformant having the recombinant vector.

50 A further object of the present invention is to provide a process for preparing the G-protein coupled receptor proteins or their salts.

Still another object of the present invention is to provide a method for determining ligands to the G-protein coupled receptors.

55 Still another object of the present invention is to provide a method for screening for receptor-agonists or antagonists to the G-protein coupled receptor proteins, a kit for screening for the receptor-agonists or antagonists, the receptor-agonists or antagonists obtained by the screening and a pharmaceutical composition comprising at least one of the receptor-agonists or antagonists.

Yet another object of the present invention is to provide antibodies against the G-protein coupled receptor proteins, their partial peptides or salts.

These objects as well as other objects and advantages of the present invention will become apparent to those skilled in the art from the following description with reference to the accompanying drawings.

## BRIEF EXPLANATION OF DRAWINGS

Fig. 1 is a nucleotide sequence encoding the human G-protein coupled receptor protein (short form) of the present invention obtained in Example 1 hereinafter and its amino acid sequence deduced from the nucleotide sequence.

Fig. 2 is a graph illustrating hydrophobic plotting of the human G-protein coupled receptor protein (short form) of the present invention prepared based on the amino acid sequence of Fig. 1. The parts represented by 1 to 7 are hydrophobic domains.

Fig. 3 is a nucleotide sequence encoding the human G-protein coupled receptor protein (long form) of the present invention obtained in Example 1 hereinafter and its amino acid sequence deduced from the nucleotide sequence.

Fig. 4 is a graph illustrating hydrophobic plotting of the human G-protein coupled receptor protein (long form) of the present invention prepared based on the amino acid sequence of Fig. 1. The parts represented by 1 to 7 are hydrophobic domains.

Fig. 5 illustrates the results of northern hybridization for examining expression levels of mRNA encoding the human G-protein coupled receptor protein of the present invention in various human tissues. The value (kb) represents the size of the RNA molecular weight marker.

## SUMMARY OF THE INVENTION

As a result of an intensive study, the present inventors have succeeded in isolation of cDNAs encoding two kinds of G-protein coupled receptor proteins derived from human fetal brain and human adult brain based on two kinds of publicly available EST information registered with a data base whose functions are unknown, and have succeeded in analysis of the entire nucleotide sequences. When their amino acid sequences have been deduced from the nucleotide sequences, the first to the seventh transmembrane domains have been confirmed on hydrophobic plotting. Thus, the proteins encoded by these cDNAs have been confirmed to be 7-transmembrane type G-protein coupled receptor proteins. The present invention has been completed based on these findings.

That is, according to the present invention, there are provided:

- (1) A G-protein coupled receptor protein which comprises the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 1, or its salt;
- (2) The G-protein coupled receptor protein of the above (1) which comprises the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 2, or its salt;
- (3) A partial peptide of the G-protein coupled receptor protein of the above (1) or its salt;
- (4) An isolated DNA comprising DNA having a nucleotide sequence encoding the G-protein coupled receptor protein of the above (1);
- (5) The isolated DNA of the above (4) having the nucleotide sequence represented by SEQ ID NO: 3;
- (6) The isolated DNA of the above (4) having the nucleotide sequence represented by SEQ ID NO: 4;
- (7) A recombinant vector comprising the DNA of the above (4);
- (8) A transformant comprising the recombinant vector of the above (7);
- (9) A process for preparing the G-protein coupled receptor protein of the above (1) or its salt which comprises culturing the transformant of the above (8) to form the G-protein coupled receptor protein;
- (10) A method for determining a ligand to the G-protein coupled receptor protein of the above (1) or its salt which comprises bringing the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt into contact with a test compound;
- (11) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises comparing (i) ligand binding upon bringing the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt into contact with the ligand, and (ii) that upon bringing the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt into contact with the ligand and a test compound;
- (12) A kit for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises as an essential component the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt;
- (13) The compounds which alter ligand binding to the G-protein coupled receptor protein of the above (1) or its salt obtained by the screening method of the above (11) or the kit of the above (12), or their salts; and
- (14) An antibody against the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt.

More specifically, the present invention provides:

(15) The G-protein coupled receptor protein of the above (1) or its salt, wherein the protein comprises the amino acid sequence represented by SEQ ID NO: 1, a variant of the amino acid sequence represented by SEQ ID NO: 1 having a deletion of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids, a variant of the amino acid sequence represented by SEQ ID NO: 1 having an addition of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids, or a variant of the amino acid sequence represented by SEQ ID NO: 1 having a substitution of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids;

(16) The G-protein coupled receptor protein of the above (2) or its salt, wherein the protein comprises the amino acid sequence represented by SEQ ID NO: 2, a variant of the amino acid sequence represented by SEQ ID NO: 2 having a deletion of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids, a variant of the amino acid sequence represented by SEQ ID NO: 2 having an addition of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids, or a variant of the amino acid sequence represented by SEQ ID NO: 2 having a substitution of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids;

(17) The method for determining a ligand of the above (10), wherein the ligand is angiotensin, bombesin, cannabinoid, cholecystokinin, glutathione, melatonin, neuropeptide Y, opiod, purine, vasopressin, oxytocin, PAC-AP, secretin, glucagon, calcitonin, adrenomedullin, somatostatin, GHRH, CRF, ACTH, GRP, PTH, VIP (vasoactive intestinal and related polypeptides), dopamine, melin, amylin, bradykinin, CGRP (calcitonin gene related peptide), leukotriene, pancreastatin, prostaglandin, thromboxane, adenosine, adrenalin,  $\alpha$  or  $\beta$ -chemokine (e.g., L-8, GRO $\alpha$ , GRO $\beta$ , GRO $\gamma$ , NAP-2, ENA-78, PFA, IP10, GCP-2, MCP-1, HC14, MCP-3, I-305, MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, etc.), endothelin, enterogastin, histamine, neurotensin, TRH, pancreatic polypeptides or galanin;

(18) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein or its salt of the above (1), or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to the G-protein coupled receptor protein of the above (1) or its salt or the partial peptide of the above (3) or its salt upon bringing the protein of the above (1), the partial peptide of the above (3) or a salt thereof into contact with the labeled ligand and a test compound;

(19) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to cells containing the G-protein coupled receptor protein of the above (1) upon bringing the labeled ligand into contact with the cell membrane fraction, and (ii) that upon bringing the labeled ligand into contact with the cells with (i) that upon bringing the labeled ligand and a test compound into contact with the cells;

(20) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to a membrane fraction of cells containing the G-protein coupled receptor protein of the above (1) upon bringing the labeled ligand into contact with the cell membrane fraction, and (ii) that upon bringing the labeled ligand and a test compound into contact with the cell membrane fraction;

(21) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to the G-protein coupled receptor protein expressed on the cell membrane of the transformant of the above (8) by culturing the transformant upon bringing the labeled ligand into contact with the expressed G-protein coupled receptor protein, and (ii) that upon bringing the labeled ligand and a test compound into contact with the expressed G-protein coupled receptor protein;

(22) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises measuring and comparing (i) a cell stimulation activity mediated by the G-protein coupled receptor protein upon bringing a compound which activates the G-protein coupled receptor protein of the above (1) or its salt into contact with cells containing the G-protein coupled receptor protein of the above (1), and (ii) that upon bringing the compound which activates the G-protein coupled receptor protein or its salt and a test compound into contact with the cells;

(23) A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or its salt, or their salts which comprises measuring and comparing (i) a cell stimulation activity mediated by the G-protein coupled receptor protein upon bringing a compound which activates the G-protein coupled receptor protein of the above (1) or its salt into contact with the G-protein coupled receptor protein expressed on the cell membrane of the transformant of the above (8) by culturing the transformant, and (ii) that upon bringing the compound which activates the G-protein coupled receptor protein or its salt and a test compound into contact with the G-protein coupled receptor protein expressed on the cell membrane;

(24) The method for screening of the above (22) or (23), wherein the compound which activates the G-protein coupled receptor protein of the above (1) is angiotensin, bombesin, cannabinoid, cholecystokinin, glutamine, se-

ratonin, melatonin, neuropeptide Y, opiod, purine, vasopressin, oxytocin, PACAP, secinin, glucagon, calcitonin, adrenomedullin, somatostatin, GHRH, CRF, ACTH, GHRP, PTH, VIP (vasoactive intestinal and related polypeptides), dopamine, melatonin, amylin, bradykinin, CGRP (calcitonin gene related proteins), leukotriene, pancreaticatin, histidine, prostaglandin, thromboxane, adenosine, adrenalin,  $\alpha$  or  $\beta$ -adrelinic (e.g., IL-8, GHRQ, GHRQ, GHRQ, MAP-2, ENA-7B, P-PE4, IP10, GCR-2, MCP-1, HCl4, MCP-3, 1309, MIP-1 $\beta$ , RANTES, etc.), endothelin, antro-5, gastrin, histamine, neuropeptides, TSH, pancreatic polypeptides or gallamine;

(25) Compounds which alter binding of a ligand to the G-protein coupled receptor protein of the above (1) or (24), or their salts; salt obtained in any one of the method of screening of the above (11) and (18) or (24), or their salts;

(26) A pharmaceutical composition comprising the compound of the above (25) or its salt;

(27) The screening kit of the above (12) comprising cells containing the G-protein coupled receptor protein of the above (11);

(28) The screening kit of the above (12) comprising a membrane fraction of whole cells containing the receptor protein of the above (1);

(29) The compound which alter ligand binding to the G-protein coupled receptor protein of the above (1) or its salt obtained by using the screening kit of the above (12), (27) or (28); or its salt.

(30) A pharmaceutical composition comprising the compound of the above (29) or its salt, and  
(31) A method for determining the G-protein coupled receptor protein of the above (1), or its salt or the partial peptide of the above (3) or its salt which comprises bringing the antibody of the above (14) into contact with the peptide of the above (3) or its salt, the partial peptide of the above (3) or a salt thereof.  
G-protein coupled receptor protein of the above (1), the partial peptide of the above (3) or a salt thereof.

### DETAILED DESCRIPTION OF THE INVENTION

The G-protein coupled receptor protein of the present invention, (hereinafter sometimes abbreviated to "receptor protein") is the receptor protein which has the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 1 (Fig. 1) and it may be the receptor protein having the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 2 (Fig. 3). The amino acid sequence of SEQ ID NO. 2 is a variant of the amino acid sequence represented by SEQ ID NO. 1 having an addition of 61 amino acids at the N-terminal end and SEQ ID NO. 1.

The receptor protein of the present invention may be any peptide derived from any cells of a human being and another mammal (e.g., guinea pig, rat, mouse, chicken, rabbit, pig, sheep, cattle, monkey, etc.), for example, specific cells, nerve cell, glia cell of perineur, marrow cell, mesangial cell, Langerhans cell, epidermic cell, epithelial cell, endothelial cell, fibroblast, fibrocyte, muscular cell, fat cell, immunocyte (e.g., macrophage, T cell, B cell, natural killer cell, mast cell, neutrophil, basophil, eosinophilic leukocyte, monocyte, etc.), megakaryocyte, myeloid cell, dendritic cell, osteocyte, osteoblast, osteoclast, mammary gland cell, hepatocyte, or intestinal cells or precursor cells, stem cells or cancer cells thereof and the like, and any tissues containing such cells, for example, brain, various parts of brain (e.g., olfactory bulb, amygdala, cerebral basal ganglia, hippocampus, thalamus, hypothalamus, caudate nucleus, subthalamic nucleus, cerebellar cortex, medulla, cerebellum, occipital pole, frontal lobe, pituitary, caudate nucleus, corpus callosum, substantia nigra), spinal cord, pituitary, stomach, pancreas, kidney, liver, genital organs, thyroid gland, gallbladder, bone marrow, adrenal gland, skin, muscle, lung, digestive tract, blood vessel, heart, thymus, spleen, submandibular gland, peripheral blood, peripheral blood leukocyte, intestinal tract, prostate, testicle, ovary, placenta, uterus, bone marrow, epithelium, small intestine, large intestine, skeletal muscle and the like, in particular, brain and various parts of brain. The peptide may be a synthetic one.

The wording 'the same or substantially the same as the amino acid sequence as that represented by SEQ ID NO. 1 or SEQ ID NO. 2' includes any protein which has at least about 70% homology, preferably at least about 80%, 1 or SEQ ID NO. 2, to the amino acid sequence represented by SEQ ID NO. 1 or SEQ ID NO. 2, and having substantially the same activity as that of the receptor protein comprising the amino acid sequence represented by SEQ ID NO. 1 or SEQ ID NO. 2.

Examples of substantially the same activity include ligand binding activity signal information transmission activity and the like. The wording "substantially the same" means that the natures of their activities are equal to one another. Therefore, quantitative factors such as degrees of ligand binding activity and signal information transmission activity may differ from one another.

Further, the receptor protein of the present invention may be a protein comprising a variant of the amino acid sequence represented by SEQ ID NO. 2 having a deletion of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids, a variant of the amino acid sequence represented by SEQ ID NO. 2 having an addition of one or more, preferably about 1 to about 10, more preferably about 1 to about 10 amino acids, or a variant of the amino acid sequence represented by SEQ ID NO. 2 having a substitution of one or more, preferably about 1 to about 30, more preferably about 1 to about 10 amino acids.

More specifically, the receptor protein of the present invention includes, for example, the receptor protein compris-

ing the amino acid sequence represented by SEQ ID: NO 1 derived from a human brain, or the G-protein coupled receptor protein comprising the amino acid sequence represented by SEQ ID: NO 2 derived from a human brain.

Furthermore, examples of the receptor protein of the present invention include variants of the above receptor protein, wherein the amino group of the N-terminal methionine residue of the above receptor protein is protected with a protecting group (e.g., -acyl group having 1 to 6 carbon atoms such as tomyl group, acetyl group, etc.); the N-terminal amino group (e.g., -acyl group having 1 to 6 carbon atoms such as tomyl group, acetyl group, etc.); the N-terminal amino group (e.g., -OH, -COOH, amino group, imidazole group, indole group, guanidino group, etc.) on the side chain of a substituent (e.g., -OH, -COOH, amino group, imidazole group, indole group, guanidino group, etc.) of an amino acid in the molecule of the above receptor protein is protected with a suitable protecting group (e.g., -acyl group having 1 to 6 carbon atoms such as tomyl group, acetyl group, etc.); or conjugated proteins of the above receptor protein having 1 to 6 carbon atoms such as tomyl group, acetyl group, etc.

The receptor protein of the present invention is represented by a conventional nomenclature in peptide art. That is, the left hand end (amino terminal) is the N-terminal and the right hand end (carboxyl terminal) is the C-terminal. And, in the receptor protein of the present invention, a representative example thereof being the receptor protein comprising the amino acid sequence represented by SEQ ID NO. 1 or SEQ ID NO. 2, normally, the amino acid is carboxyl group (-COOH) or carboxylate (-COO<sup>-</sup>), but the C-terminal may be the amide (-CONH<sub>2</sub>) or an ester (-COOR). Examples of R of the ester group include an alkyl group having 1 to 6 carbon atoms such as cyclopentyl, cyclohexyl, etc., an aryl group having 6 to 10 carbon atoms such as phenyl,  $\alpha$ -naphthyl, etc., an allyl group having 7 to 14 carbon atoms such as phenyl-C<sub>3</sub>-4,  $\alpha$ -naphthyl-C<sub>3</sub>-4, alkyl group (e.g.,  $\alpha$ -naphthylmethyl, etc.) and the like. In addition, preferably, carboxyl group or the like which is used widely as an ester for oral administration can also be used.

When the receptor protein of the present invention has a carboxy group (or carboxylic acid) as a protein secretion signal, the amino acid sequence of the amino acid residue immediately preceding the carboxy group is preferably an esterifiable amino acid. When the receptor protein of the present invention has a carboxy group (or carboxylic acid) as a protein secretion signal, the amino acid sequence of the amino acid residue immediately preceding the carboxy group is preferably an esterifiable amino acid. When the receptor protein of the present invention has a carboxy group (or carboxylic acid) as a protein secretion signal, the amino acid sequence of the amino acid residue immediately preceding the carboxy group is preferably an esterifiable amino acid.

As the salt of the receptor protein of the present invention, in particular, a physiologically acceptable acid addition salt is preferred. Examples of the salt include those with inorganic acids (e.g., hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid, etc.) and those with organic acids (e.g., acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, methanesulfonic acid, benzenesulfonic acid, etc.).

The receptor protein or its salt of the present invention can be prepared from the above-described cells and tissues of a human being or another mammal by a per se known purification method of proteins. Alternatively, the receptor of a human being or another mammal can be prepared by culturing a transformation containing DNA encoding the protein or its salt of the present invention according to a peptide synthesis method as described hereinafter.

When it is produced from cells or tissues of a human being or another mammal, the cells or tissues are homogenized and then extracted with, for example, an acid. The extract can be purified and isolated by combining chromatography (see such as reverse phase chromatography, ion exchange chromatography and the like,

As the partial peptide of the receptor protein (hereinafter sometimes abbreviated to 'partial peptide') of the present invention, for example, a part of the receptor protein molecule of the present invention which is exposed to outside of a cell membrane or the like can be used. Specifically, the partial peptide of the receptor protein having the amino acid sequence represented by SEQ ID NO: 1 is a peptide containing the parts which have been analyzed to the extracellular sequence represented by SEQ ID NO: 1 in the hydrophobic plotting analysis as shown by Fig. 2. The partial peptide of the domains (hydrophobic domains) in the hydrophobic plotting analysis as shown by Fig. 2, that is, the partial peptide of the receptor protein having the amino acid sequence represented by SEQ ID NO: 2 is that containing the parts which have been analyzed to be extracellular domains (hydrophobic domains) in the hydrophobic plotting analysis as shown by Fig. 4. A peptide containing a hydrophobic domain part can be suitably used. In addition, the peptide may contain each domain separately or plural domains together.

Examples of the partial peptide include those having the amino acid sequences of the 78th to 130th amino acids of the 193rd to 204th amino acids, the 274th to the 307th amino acids, and the 397th to 398th amino acids of the amino acid sequence represented by SEQ ID NO. 1 as well as amino acid sequences of the 133th to 191st amino acids, the 254th to 265th amino acids, the 335th to 368th and the 448th to 459th amino acids of the amino acid sequence represented by SEQ ID NO. 2.

Further, the partial peptide of the receptor protein of the present invention includes variants of the above partial peptide, wherein the amino group of N-terminal methionine residue of the above receptor protein is protected with a protecting group (e.g., acyl group having 1 to 6 carbon atoms such as formyl group, acetyl group, etc.), the N-terminal region of the above receptor protein is cleaved in a living body and the glutaryl group formed is pyroglutaminated, or a substituent (e.g., -OH, -COOH, amino group, indazole group, imide group, etc.) on the side chain carbon atom of an amino acid in the molecule of the above receptor protein is protected with a suitable protecting group (e.g., acyl group having 1 to 6 carbon atoms such as formyl group, acetyl group, etc.), or conjugated peptides of the above partial peptide such as glycopeptides having sugar chains.

Normally, the C-terminal of the partial peptide of the present invention is a carboxyl group (-COOH) or carboxylate (-COO<sup>-</sup>) and, like the receptor protein of the present invention, the C-terminal may be the amide or ester. When the partial peptide of the present invention has a carboxyl group (or carboxylate) at a position other than the C-terminal, it may be amidated or esterified and such amide or ester is also included in the scope of the partial peptide of the present invention. The ester group may be the same group as that described with respect to the above C-terminal of the receptor protein.

As the salt of the partial peptide of the present invention, in particular, a physiologically acceptable acid addition salt is preferred. Examples of the salt include those with inorganic acids (e.g., hydrochloric acid, phosphoric acid, hydrobromic acid, sulfuric acid, etc.) and those with organic acids (e.g., acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, benzoesulfonic acid, benzenesulfonic acid, etc.).

The partial peptide or its salt of the present invention can be prepared according to a per se known peptide synthesis method or by cleaving the receptor protein of the present invention with a suitable peptidase.

As the peptide synthesis method, for example, any of solid phase synthesis and liquid phase synthesis can be employed. That is, the objective peptide can be produced by condensing a partial peptide or amino acid sequence which can compose of the partial peptide of the present invention with the remaining part and deprotecting a protecting group, if any. Conventional condensing methods and deprotecting methods can be employed and they are described by, for example, M. Bodanszky and M.A. Ordeit, *Peptide Synthesis*, Interscience Publishers, New York (1966); Schroeder and Liebeck, *The Peptide, Academic Press*, New York (1965); Nobuo Izumi et al., *Fundamental and Experiment of Peptide Synthesis*, Maruzen (1975); Haruaki Yazima and Syunpei Skakibara, *Biochemistry Experiment Lecture*, Protein Chemistry IV, 205 (1977); Haruaki Yazima, *Second Series Drug Development Vol. 14*, Peptide Synthesis, Hirokawa Shoten.

After completion of the reaction, the partial peptide of the present invention can be purified and isolated by combining conventional purification methods such as solvent extraction, distillation, column chromatography, liquid chromatography, recrystallization and the like. In case the partial peptide thus obtained is a free peptide, it can be converted into its appropriate salt according to a known method. On the other hand, the peptide obtained is in the form of a salt, it can be converted into the corresponding free peptide.

The DNA encoding the receptor protein of the present invention may be any DNA in so far as it contains the nucleotide sequence encoding the above-described receptor protein of the present invention. The DNA may be any of genomic DNA, genomic DNA library, cDNA derived from the above-described cells and tissues, cDNA library derived from the above-described cells and tissues and synthetic DNA. The vector to be used for the library may be any of bacteriophage, plasmid, cosmid, phagemid and the like. In addition, the DNA can be amplified by reverse transcriptase polymerase chain reaction (hereinafter abbreviated to RT-PCR) with a mRNA fraction prepared from the above-described cells and tissues.

Specifically, the DNA encoding the receptor protein having the same or substantially the same amino acid sequence represented by SEQ ID NO: 1 of the present invention may be, for example, DNA having the nucleotide sequence represented by SEQ ID NO: 3 or any DNA having a nucleotide sequence hybridizable to the nucleotide sequence represented by SEQ ID NO: 3 under high stringent conditions and encoding a receptor protein which has the same activities, i.e., ligand binding activity, signal information transmission activity and the like as those of the receptor protein having the amino acid sequence represented by SEQ ID NO: 1. Examples of the hybridizable DNA include DNA having at least about 70% to about 80% homology, preferably, at least about 90% homology, more preferably, at least about 95% homology to the nucleotide sequence represented by SEQ ID NO: 3.

More specifically, as the DNA encoding the receptor protein containing the amino acid sequence represented by SEQ ID NO: 1, the DNA having the nucleotide sequence represented by SEQ ID NO: 3 or the like can be used.

The DNA encoding the receptor protein having the same or substantially the same amino acid sequence represented by SEQ ID NO: 2 of the present invention may be, for example, DNA having the nucleotide sequence represented by SEQ ID NO: 4 or any DNA having a nucleotide sequence hybridizable to the nucleotide sequence represented by SEQ ID NO: 4 under high stringent conditions and encoding a receptor protein which has the same activities, i.e., ligand binding activity, signal information transmission activity and the like as those of the receptor protein having the amino acid sequence represented by SEQ ID NO: 2. Examples of the hybridizable DNA include DNA having at least about 70% to about 80% homology, preferably, at least about 90% homology, more preferably, at least about 95% homology to the nucleotide sequence represented by SEQ ID NO: 4.

More specifically, as the DNA encoding the receptor protein containing the amino acid sequence represented by SEQ ID NO: 2, the DNA having the nucleotide sequence represented by SEQ ID NO: 4 or the like can be used. The nucleotide sequence represented by SEQ ID NO: 4 is a variant of the nucleotide sequence of SEQ ID NO: 3 having an addition of 183 bases at its 5'-terminal.

Hybridization can be carried out by a per se known method or its modification, for example, under high stringent conditions.

The high stringent conditions used herein are, for example, those of sodium concentration at about 19 mM to about 40 mM, preferably about 19 mM to about 20 mM and a temperature at about 50°C to about 70°C, preferably about 60°C to about 65°C. In particular, hybridization conditions of sodium concentration at about 19 mM and a temperature at about 65°C are most preferred.

The DNA encoding the partial peptide of the present invention may be any DNA in so far as it contains the nucleotide sequence encoding the above-described partial peptide of the present invention. The DNA may be any of genomic DNA, genomic DNA library, cDNA derived from the above-described cells and tissues, cDNA library derived from the above-described cells and tissues and synthetic DNA. The vector to be used for the library may be any of bacteriophage, plasmid, cosmid, phagemid and the like. In addition, the DNA can be amplified by reverse transcriptase polymerase chain reaction (hereinafter abbreviated to RT-PCR) with a mRNA fraction prepared from the above-described cells and tissues.

Specifically, the DNA encoding the partial peptide of the receptor protein having the same or substantially the same amino acid sequence represented by SEQ ID NO: 1 of the present invention may be, for example, DNA having the nucleotide sequence represented by SEQ ID NO: 3 or any DNA having a nucleotide sequence hybridizable to the nucleotide sequence represented by SEQ ID NO: 3 under high stringent conditions and encoding a receptor protein which has the same activities, i.e., ligand binding activity, signal information transmission activity and the like as those of the receptor protein having the amino acid sequence represented by SEQ ID NO: 1. Examples of the hybridizable DNA include DNA having at least about 70% to about 80% homology, preferably, at least about 90% homology, more preferably, at least about 95% homology to the nucleotide sequence represented by SEQ ID NO: 3.

For example, as the DNA encoding the partial peptide of the receptor protein containing the amino acid sequence represented by SEQ ID NO: 1, the DNA having the nucleotide sequence represented by SEQ ID NO: 3 or the like can be used. More specifically, as the DNA encoding the partial peptide having the amino acid sequences of the 78th to 130th amino acids, the 139th to 204th amino acids, the 274th to the 307th amino acids or the 387th to 398th amino acids of the amino acid sequence represented by SEQ ID NO: 1, the DNA having the nucleotide sequence of the 232nd to 390th bases, the 577th to 612th bases, the 620th to 921st bases or the 1159th to 1194th bases of the nucleotide sequence represented by SEQ ID NO: 3 can be used.

The DNA encoding the partial peptide of the receptor protein having the same or substantially the same amino acid sequence represented by SEQ ID NO: 2 of the present invention may be, for example, DNA having the nucleotide sequence represented by SEQ ID NO: 4 or any DNA having a nucleotide sequence hybridizable to the nucleotide sequence represented by SEQ ID NO: 4 under high stringent conditions and encoding a receptor protein which has the same activities, i.e., ligand binding activity, signal information transmission activity and the like as those of the receptor protein having the amino acid sequence represented by SEQ ID NO: 2. Examples of the hybridizable DNA include DNA having at least about 70% to about 80% homology, preferably, at least about 90% homology, more preferably, at least about 95% homology to the nucleotide sequence represented by SEQ ID NO: 4.

For example, as the DNA encoding the partial peptide of the receptor protein containing the amino acid sequence represented by SEQ ID NO: 2, the DNA having the nucleotide sequence represented by SEQ ID NO: 4 or the like can be used. More specifically, as the DNA encoding the partial peptide having the amino acid sequences of the 139th to 191st amino acids, the 254th to 265th amino acids, the 335th to the 368th amino acids or the 448th to 459th amino acids of the amino acid sequence represented by SEQ ID NO: 2, the DNA having the nucleotide sequence of the 415th to 573rd bases, the 760th to 795th bases, the 1003rd to 1104th bases or the 1342nd to 1377th bases of the nucleotide sequence represented by SEQ ID NO: 4 can be used.

Hybridization can be carried out by a per se known method or its modification, for example, under high stringent conditions.

As described above, the high stringent conditions used herein are, for example, those of sodium concentration at about 19 mM to about 40 mM, preferably about 19 mM to about 20 mM, and a temperature at about 50°C to about 70°C, preferably about 60°C to about 65°C. In particular, hybridization conditions of sodium concentration at about 19 mM and a temperature at about 65°C are most preferred.

As the means for cloning the DNA encoding the entire receptor protein of the present invention, there is amplification by PCR using synthetic DNA primers containing partial nucleotide sequences of the receptor protein of the present invention. Alternatively, DNA integrated into a suitable vector is selected by hybridization with labeled DNA fragment or a synthetic DNA encoding a part or entire region of the receptor protein of the present invention. The hybridization is carried out, for example, according to the method described in Molecular Cloning 2nd Ed., J. Sambrook et al., Cold Spring Harbor Lab. Press, (1989). A commercially available library can be used according to the direction of the attached manufacturer's protocol.

Conversion of the nucleotide sequence of DNA can be carried out according to a per se known method such as Gapped duplex method or Kunkel method or its modification by using a known kit, Muta<sup>TM</sup>-K of Mutan<sup>TM</sup>-K (both Takara Shuzo Co., Ltd., TM represents trademark). The nucleotide sequence represented by SEQ ID NO: 1 can be produced by deleting 183 bases from the 5'-terminal of the nucleotide sequence represented by SEQ ID NO: 2.



The cloned DNA encoding the receptor protein can be used as such according to a particular purpose. Alternatively, it desired, it can be used after digestion with one or more restriction enzymes or a linker can be added. The DNA may have the codon, ATG, as a translation initiation codon at its 5' terminal side and the codon, TAA, TGA or TAG as a translation termination codon at its 3' terminal side. These translation initiation and termination codons can be added by using a suitable synthetic DNA adapter.

The expression vector of the receptor protein of the present invention can be prepared, for example, by (a) cutting the desired DNA fragment from the DNA encoding the receptor protein of the present invention and (b) joining the DNA fragment to a suitable expression vector at the downstream from a promoter in the vector.

Examples of the vector include plasmids derived from *E. coli* (e.g., pBR322, pBR325, pUC12, pUC13), plasmids derived from *Bacillus subtilis* (e.g., pUB110, pTP5, pC194), plasmids derived from yeast (e.g., pSH19, pSH15), bacteriophages such as  $\lambda$  phage, etc., animal viruses such as reovirus, vaccinia virus, baculovirus, etc. as well as pA1-11, pX11, pBrCMV, pBrCMV, pCDNA/Neo, etc.

The promoter used in the present invention may be any promoter in so far as it matches with a host to be used for gene expression. In case of using animal cells as the host, examples of the promoter include SP6 promoter, SV40 promoter, LTR promoter, CMV promoter, HSV-TK promoter, etc. Among them, CMV promoter or SP6 promoter is preferred. In case of using bacteria of the genus *Escherichia* as the host, preferred examples of the promoter include lac promoter, lacZ promoter,  $\lambda$ P<sub>1</sub> promoter, lpp promoter, etc. In case of using bacteria of the genus *Bacillus* as the host, preferred examples of the promoter include PHOS promoter, PGK promoter, GAP promoter, ADH promoter, etc. In case of using insect cells as the host, preferred examples of the promoter include polyhedrin promoter, P10 promoter, etc.

In addition to the above, optionally, the expression vector may further contain enhancer, splicing signal, poly A addition signal, selection marker, SV40 replication origin (hereinafter sometimes abbreviated to SV40 ori) etc. Examples of the selection marker include dihydrodolate reductase (hereinafter sometimes abbreviated to dhfr) gene (neomycin resistance), ampicillin resistant gene (hereinafter sometimes abbreviated to Amp<sup>r</sup>), neomycin resistant gene (hereinafter sometimes abbreviated to Neo, G418 resistance), etc. In particular, when CHO (dhfr<sup>-</sup>) cell is used together with DHFR gene as the selection marker, selection can also be carried out by using a thymidine free medium.

If necessary, a signal sequence which matches with a host is added to the N-terminal side of the receptor protein of the present invention. As the signal sequence, there may be mentioned alkaline phosphatase signal sequence, OmpA signal sequence, etc. in case of using bacteria of the genus *Escherichia* as the host, a-amylase signal sequence, subtilisin signal sequence, etc. in case of using bacteria of the genus *Bacillus* as the host, mating factor, a signal sequence, invertase signal sequence, etc. in case of using yeast as the host, insulin signal sequence,  $\alpha$ -interferon signal sequence, antibody molecule signal sequence, etc. in case of using animal cells as the host, respectively.

The DNA encoding receptor protein of the present invention thus constructed can be employed to transform the host.

As the host, for example, there may be mentioned bacteria of the genus *Escherichia*, bacterioid of the genus *Bacillus*, yeast, insect cells, insects and animal cells, etc.

Specific examples of bacteria of the genus *Escherichia* include *Escherichia coli* K12 DH1 (Proc. Natl. Acad. Sci. USA, 80, 180 (1983)), JM103 [Nucleic Acids Research, 9, 309 (1981)], JAZ21 [Journal of Molecular Biology, 120, 517 (1978)], HB101 [Journal of Molecular Biology, 41, 459 (1989)], C600 [Genetics, 99, 440 (1984)], etc.

Examples of bacteria of the genus *Bacillus* include *Bacillus subtilis* M114 [Gene, 24, 255 (1983)], 207-21 [Journal of Microbiology, 95, 87 (1984)], etc.

Examples of yeast include *Saccharomyces cerevisiae* AH22, AH22-NA87-11A, DKD-50, 208-12, etc.

Examples of insect cells include *Spodoptera frugiperda* cell (Sf cell), MG1 cell derived from mid-intestine of *T. chlopius* p1, High Five™ cell derived from egg of *T. chlopius* p1, cells derived from *Mamestra brassicae*, cells derived from *Estigmene acrea*, etc. for the virus, AcNPV, and Bombyx mori N cell (BmN cell), etc. for the virus, BmNPV. As the Sf cell, for example, Sf9 cell (ATCC CRL 1711) and Sf21 cell described by Vaughn, J. L., in *Viro*, 13, 213-217 (1977) can be used.

As the insect, for example, a larva of *Bombyx mori* can be used [Nakada et al., Nature, 315, 592 (1985)].

Examples of animal cells include monkey cell COS-7, Vero, Chinese hamster cell CHO, DHFR gene deficient Chinese hamster cell CHO (dhfr<sup>-</sup> CHO cell), mouse L cell, mouse AIT-20, mouse myeloma cell, rat GH 4, human FL cell, etc.

Transformation of bacteria of the genus *Escherichia* is carried out, for example, according to the method described in Proc. Natl. Acad. Sci. USA, 89, 2110 (1972) or Gene, 17, 107 (1982).

Transformation of bacteria of the genus *Bacillus* is carried out, for example, according to the method described in Molecular & General Genetics, 188, 111 (1979).

Transformation of yeast is carried out, for example, the method described in Proc. Natl. Acad. Sci. USA, 75, 1929 (1978).

Transformation of insect cells or insect is carried out, for example, according to the method described in Bio Technology, 5, 47-55 (1988).

Transformation of animal cells is carried out, for example, according to the method described in Virology, 52, 456 (1973).

Thus, the transformant transformed with the expression vector containing the DNA encoding the G-protein coupled receptor protein can be obtained.

In case of the bacterial host of the genus *Escherichia* or *Bacillus*, the transformant can be suitably cultivated in a liquid culture medium and materials required for growth of the transformant such as carbon sources, nitrogen sources, inorganic materials, etc. are added to the medium. Examples of the carbon sources include glucose, dextrin, soluble starch, sucrose, etc. The nitrogen sources include, for example, inorganic or organic materials such as ammonium salts, nitrate salts, casein, meat extract, soybean meal, potato extract, etc. The inorganic materials include, for example, calcium chloride, sodium dihydrogen phosphate, magnesium chloride, etc. In addition, yeast, vitamins, growth promoting factors etc. can be added. Preferably, the medium is adjusted to pH about 5 to about 8. Preferably, the medium for cultivating the bacteria of the genus *Escherichia* is, for example, M9 medium containing glucose and Casamino Acids (Miller, Journal of Experiments in Molecular Genetics, 431-433, Cold Spring Harbor Laboratory, New York, 1972). If necessary, in order to activate the promoter efficiently, for example, an agent such as 3 $\beta$ -indolyl acryic acid can be added to the medium.

In case of the bacterial host of the genus *Escherichia*, normally, the transformant is cultivated at about 15°C to about 43°C for about 3 hours to about 24 hours. If necessary, the culture can be aerated or stirred.

In case of the bacterial host of the genus *Bacillus*, normally, the transformant is cultivated at about 30°C to about 40°C for about 6 hours to about 24 hours. If necessary, the culture can be aerated or stirred.

In case of the yeast host, the transformant is cultivated in, for example, Burohoder's minimal medium [Bostian, K. L. et al., Proc. Natl. Acad. Sci. USA, 77, 4505 (1980)] and SD medium containing 0.5% Casamino Acids [Miller, G. A., Proc. Natl. Acad. Sci. USA, 81, 5330 (1984)]. Preferably, the medium is adjusted to pH about 5 to about 8. Normally, the transformant is cultivated at about 20°C to about 35°C for about 24 hours to about 72 hours. If necessary, the culture can be aerated or stirred.

In case of the insect cell host or insect host, the transformant is cultivated in, for example, Grace's Insect Medium [Grace, T. C., Nature, 195, 788 (1962)] to which an appropriate additive such as inactivated 10% bovine serum is added. Preferably, the medium is adjusted to pH about 6.2 to about 6.4. Normally, the transformant is cultivated at about 27°C for about 3 days to about 5 days and, if necessary, the culture can be aerated or stirred.

In case of the animal cell host, the transformant is cultivated in, for example, MEM medium containing about 5% to about 20% fetal bovine serum [Science, 122, 501 (1952)], DMEM medium [Virology, 8, 398 (1959)], RPMI 1640 medium [The Journal of the American Medical Association, 199, 519 (1957)], 199 medium [Proceeding of the Society for the Biological Medicine, 73, 1 (1950)], etc. Preferably, the medium is adjusted to pH about 6 to about 8. Normally, the transformant is cultivated at about 30°C to about 40°C for about 15 hours to about 60 hours and, if necessary, the culture can be aerated or stirred.

As described hereinabove, the G-protein coupled receptor protein of the present invention can be produced at the cell membrane of the transformant.

Separation and purification of the receptor protein of the present invention from the above culture can be carried out, for example, as follows.

Extraction of the receptor protein of the present invention from the transformant culture can be carried out by an appropriate known method. For example, after cultivation, the transformant is recovered by a per se known method and suspended in a suitable buffer. Then, the transformant is disrupted by a per se known method such as ultrasonic-cation, treatment with lysozyme and/or freeze-thaw cycling, followed by separating a crude extract of the receptor protein by centrifugation, filtration, etc. The buffer may contain a protein modifier such as urea, guanidine hydrochloride or a surfactant such as Triton X-100™, etc. When the receptor protein is secreted in the culture broth, after completion of cultivation, its supernatant can be separated from the transformant cells to collect the supernatant.

Purification of the receptor protein contained in the culture supernatant thus obtained or the extract can be carried out by combining per se known separation and purification methods appropriately. As the per se known separation and purification methods, there may be mentioned a method utilizing difference in solubilities such as salting out, solvent precipitation, etc.; a method mainly utilizing difference in molecular weights such as dialysis, ultrafiltration, gel filtration, SDS-polyacrylamide gel electrophoresis, etc.; a method utilizing difference in electric charges such as ion exchange chromatography, etc.; a method utilizing difference in specific affinities such as affinity chromatography, etc.; a method utilizing difference in hydrophobic properties such as reverse phase high performance liquid chromatography, etc.; a method utilizing difference in isoelectric points such as isoelectric point electrophoresis; and the like.

When the free receptor protein is obtained, it can be converted into its salt by a per se known method or its modification. On the other hand, when the receptor protein is obtained in the form of a salt, it can be converted into the free receptor protein or a different salt by a per se known method or its modification.

The receptor protein produced by the recombinant can be treated with an appropriate protein modifying enzyme prior to or after purification to appropriately modify the protein or to partially remove a polypeptide. Examples of the protein modifying enzyme include trypsin, chymotrypsin, arginyl endopeptidase, protein kinase, glycosidase and the like.

The activity of thus-produced receptor protein or its salt of the present invention can be determined by a binding test with a labeled ligand or an enzyme immunoassay with a specific antibody.

The receptor protein, its partial peptide or their salts, and DNAs encoding them of the present invention can be employed (a) in a method for determination of a ligand to the receptor protein of the present invention, (b) for obtaining an antibody or an antiserum, (c) for construction of a expression system of the recombinant receptor protein, (d) for development of an receptor binding assay system and screening for candidate compounds for drugs using the expression system, (e) for practice of drug design based on comparison with structurally analogous ligands and receptors, (f) as reagents for preparation of probes to be used in gene diagnosis, PCR primers, etc., (g) as drugs for gene prophylaxis and therapy, and the like.

In particular, screening for agonists or antagonists to the G-protein coupled receptor protein which are specific to a human being and another mammal can be carried out by using a receptor binding assay system utilizing an expression system of the recombinant G-protein coupled receptor protein of the present invention, and the agonists and antagonists can be used as prophylactic and therapeutic drugs for various diseases.

More specific description of the use of the receptor protein, its partial peptide or their salts, DNAs encoding the receptor protein or its partial peptide and antibody will be set forth below.

#### (f) Determination method of a ligand to the G-protein receptor protein

The receptor protein of the present invention or its salt, or the partial protein or its salt of the present invention is useful as a reagent for investigation or determination of a ligand to the receptor protein or its salt of the present invention.

That is, the present invention provides a method for determination of a ligand to the receptor protein of the present invention comprising bringing the receptor protein or its partial peptide of the present invention or a salt thereof into contact with a test compound.

Examples of the test compound include tissue extract, cell culture supernatant of a human being or another mammal (e.g., mouse, rat, pig, cattle, sheep, monkey, etc.), or the like, in addition to the above-described known ligands, such as angiotensin, bombesin, cannabinoid, cholecystokinin, glutamine, serotonin, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, PACAP, secretin, glucagon, calcitonin, adrenomedullin, somatostatin, GRH, CRF, ACTH, GPR, PTH, VIP (vasoactive intestinal and related polypeptides), dopamine, motilin, bradykinin, CGRP (calcitonin gene related proteins), leukotriene, pancreastatin, prostaglandin, thromboxane, adenosine, adrenalin,  $\alpha$ - $\beta$ -chemokine (e.g., IL-8, GRO $\alpha$ , GRO $\beta$ , GRO $\gamma$ , NAP-2, ENA-78, PF-4, IP10, GCP-2, MCP-1, HCl4, MCP-3, I-309, MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, etc.), endothelin, enterogastin, histamine, neurotensin, TRH, pancreatic polypeptides or galanin. For example, the tissue extract or the cell culture supernatant is added to the receptor protein of the present invention and the mixture is fractionated by measuring a cell stimulation activity, etc. to finally obtain a single ligand.

Specifically, the ligand determination method of the present invention is carried out by using the receptor protein or its partial peptide of the present invention or a salt thereof, or by constructing an expression system of the recombinant receptor protein and using a receptor binding assay system utilizing the expression system to determine a compound (e.g., peptide, protein, non-peptide compound, synthetic compound, fermentation product, etc.) showing a cell stimulation activity (e.g., an activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $\text{Ca}^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) upon binding to the receptor protein of the present invention.

The ligand determination method of the present invention is characterized by measurement of, for example, an amount of a test compound bound to the receptor protein or the partial peptide or a cell stimulation activity upon bringing the receptor protein or the partial peptide into contact with the test compound.

More specifically, the ligand determination method of the present invention is:

(a) a method for determination of a ligand to the receptor protein or its salt of the present invention which comprises labeling a test compound, and measuring an amount of the labeled test compound bound to the receptor protein, the partial peptide or their salts upon bringing the receptor protein, the partial peptide or their salts into contact with the labeled test compound;

(b) a method for determination of a ligand to the receptor protein or its salt of the present invention which comprises labeling a test compound, and measuring an amount of the labeled test compound bound to cells containing the receptor protein or a membrane fraction of the cells upon bringing the labeled test compound into contact with the cell or the cell membrane fraction;

(c) a method for determination of a ligand to the receptor protein which comprises labeling a test compound, and measuring an amount of the labeled test compound bound to the G-protein coupled receptor protein expressed on the cell membrane of a transformant containing the DNA encoding the receptor protein of the present invention by cultivating the transformant upon bringing the labeled test compound into contact with the expressed G-protein coupled receptor protein;

(d) a method for determination of a ligand to the receptor protein or its salt of the present invention which comprises measuring a cell stimulation activity (e.g., an activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $\text{Ca}^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor protein upon bringing a test compound into contact with cells containing the receptor protein;

(e) a method for determination of a ligand to the receptor protein or its salt which comprises which comprises measuring a cell stimulation activity (e.g., an activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $\text{Ca}^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor protein upon bringing a test compound into contact with the receptor protein expressed on the cell membrane of a transformant containing DNA encoding the receptor protein of the present invention by cultivating the transformant.

The receptor protein used in the ligand determination of the present invention may be any protein in so far as it contains the above-described receptor protein or partial peptide of the present invention. In particular, the receptor protein expressed in a large amount by using animal cells is suitable.

For producing the receptor protein of the present invention, the above-described expression process can be used. In particular, it is preferred to carry out expression of the DNA encoding the receptor protein in cells of mammals or insects. Normally, cDNA is used as a DNA fragment encoding the desired part of the protein, but the present invention is not necessarily limited to this. For example, a gene fragment or a synthetic DNA can also be used. For transferring a DNA fragment encoding the receptor protein of the present invention into a host animal cell and expressing it efficiently, it is preferred to integrate the DNA fragment into the downstream from a polyhelix promoter of nuclear polyomavirus (NPV) belonging to baculovirus whose host is insects, a promoter derived from SV40, retrovirus promoter, melon, allothionin promoter, human heat shock promoter, cytomegalovirus promoter, SRA promoter or the like. The quantitative and qualitative assay of expression of the receptor protein can be carried out by a per se known method. For example, the assay can be carried out according to the method described by Nambi, P. et al., J. Biol. Chem., 267, 19555-19559 (1992).

In the ligand determination of the present invention, the receptor protein, its partial peptide or their salts may be the receptor protein, its partial peptide or their salts as such which are purified according to a per se known method, or a material containing the receptor protein, its partial peptide or their salts such as cells containing the receptor protein or their membrane fractions.

In case of using the cells containing the receptor protein of the present invention, they may be immobilized with glutaraldehyde, formalin, etc. Immobilization can be carried out by a per se known method.

The cells containing the receptor protein of the present invention are host cells expressing the receptor protein of the present invention. The Examples of the host cells include *E. coli*, *Bacillus subtilis*, yeast, insect cells, animal cells and the like.

The cell membrane fraction is a fraction containing many cell membranes obtained by disrupting cells and then treated by a per se known method. The disruption of cells can be carried out, for example, using Potter-Elvehjem homogenizer, Waring blender or Polytron (Kinematica), ultrasonication, French press, etc. Fractionation of the cell membrane fraction can mainly be carried out by fractionation centrifugation, density-gradient centrifugation or the like. For example, a disrupted cell suspension is centrifuged at a low rate (500 rpm-3000 rpm) for a short period of time (normally, about 1 minute-10 minutes) and further the supernatant is centrifuged at a high rate (15000 rpm-30000 rpm) for about 30 minutes to about 2 hours to obtain a cell membrane fraction as precipitate. This cell membrane fraction contains the expressed receptor protein and many membrane components such as phospholipids, membrane proteins and the like.

The amount of the receptor protein contained in the cells or the cell membrane fraction is preferably  $10^5$  to  $10^6$  molecules, more preferably  $10^5$  to  $10^7$  molecules per one cell. As the expression level is higher, a ligand binding activity (specific activity) becomes higher, which makes not only construction of a high sensitive screening system but also determination of a large number of samples in one lot possible.

For carrying out the above ligand determination methods (a) to (c), a suitable receptor protein fraction and a labeled test compound are required.

The receptor protein fraction is preferably a naturally occurring receptor protein fraction or a recombinant receptor

protein fraction having the equivalent activity to that of the naturally occurring receptor protein. The equivalent activity of the equivalent fraction means the equivalent ligand binding activity, signal information transmission activity or the like.

The labelled (as compound) is preferably the above-exemplified ligand, such as angiotensin, bombesin, cannabinoid, cholecystokinin, glutamine, serotonine, melatonin, neuropeptide Y, opioid, purine, vasopressin, oxytocin, PACAP, secretin, glucagon, calcitonin, adrenomedullin, somatostatin, GHRH, CRF, ACTH, GRP, VIP, VIP-associated (histamine and related polypeptides), dopamine, motilin, amylin, bradykinin, CGRP (calcitonin gene related peptides), leukotrienes, prostaglandins, thromboxane, adenosine, adrenalin,  $\alpha$ - $\beta$ -chondromin (e.g., 1, 8- GPO, GPO3, GPO, pancreaticin, prostaglandin, thromboxane, adenosine, adrenalin,  $\alpha$ - $\beta$ -chondromin (e.g., 1, 8- GPO, GPO3, GPO, NAP-2, ENA-78, PF4, IP10, GPC-2, MCP-1, hC14, MCP-3, 1-303, MIP-1 $\alpha$ , MIP-1 $\beta$ , HANTES, etc.), endothelin, aneurysm, gastrin, histamine, neurotensin, TRH, acetylcholine or glutamate labelled with [P4], [124], [144], [147], [148], [149], [150], [151], [152], [153], [154], [155], [156], [157], [158], [159], [160], [161], [162], [163], [164], [165], [166], [167], [168], [169], [170], [171], [172], [173], [174], [175], [176], [177], [178], [179], [180], [181], [182], [183], [184], [185], [186], [187], [188], [189], [190], [191], [192], [193], [194], [195], [196], [197], [198], [199], [200], [201], [202], [203], [204], [205], [206], [207], [208], [209], [210], [211], [212], [213], [214], [215], [216], [217], [218], [219], [220], [221], [222], [223], [224], [225], [226], [227], [228], [229], [230], [231], [232], [233], [234], [235], [236], [237], [238], [239], [240], [241], [242], [243], [244], [245], [246], [247], [248], [249], [250], [251], [252], [253], [254], [255], [256], [257], [258], [259], [260], [261], [262], [263], [264], [265], [266], [267], [268], [269], [270], [271], [272], [273], [274], [275], [276], [277], [278], [279], [280], [281], [282], [283], [284], [285], [286], [287], [288], [289], [290], [291], [292], [293], [294], [295], [296], [297], [298], [299], [300], [301], [302], [303], [304], [305], [306], [307], [308], [309], [310], [311], [312], [313], [314], [315], [316], [317], [318], [319], [320], [321], [322], [323], [324], [325], [326], [327], [328], [329], [330], [331], [332], [333], [334], [335], [336], [337], [338], [339], [340], [341], [342], [343], [344], [345], [346], [347], [348], [349], [350], [351], [352], [353], [354], [355], [356], [357], [358], [359], [360], [361], [362], [363], [364], [365], [366], [367], [368], [369], [370], [371], [372], [373], [374], [375], [376], [377], [378], [379], [380], [381], [382], [383], [384], [385], [386], [387], [388], [389], [390], [391], [392], [393], [394], [395], [396], [397], [398], [399], [400], [401], [402], [403], [404], [405], [406], [407], [408], [409], [410], [411], [412], [413], [414], [415], [416], [417], [418], [419], [420], [421], [422], [423], [424], [425], [426], [427], [428], [429], [430], [431], [432], [433], [434], [435], [436], [437], [438], [439], [440], [441], [442], [443], [444], [445], [446], [447], [448], [449], [450], [451], [452], [453], [454], [455], [456], [457], [458], [459], [460], [461], [462], [463], [464], [465], [466], [467], [468], [469], [470], [471], [472], [473], [474], [475], [476], [477], [478], [479], [480], [481], [482], [483], [484], [485], [486], [487], [488], [489], [490], [491], [492], [493], [494], [495], [496], [497], [498], [499], [500], [501], [502], [503], [504], [505], [506], [507], [508], [509], [510], [511], [512], [513], [514], [515], [516], [517], [518], [519], [520], [521], [522], [523], [524], [525], [526], [527], [528], [529], [530], [531], [532], [533], [534], [535], [536], [537], [538], [539], [540], [541], [542], [543], [544], [545], [546], [547], [548], [549], [550], [551], [552], [553], [554], [555], [556], [557], [558], [559], [560], [561], [562], [563], [564], [565], [566], [567], [568], [569], [570], [571], [572], [573], [574], [575], [576], [577], [578], [579], [580], [581], [582], [583], [584], [585], [586], [587], [588], [589], [590], [591], [592], [593], [594], [595], [596], [597], [598], [599], [600], [601], [602], [603], [604], [605], [606], [607], [608], [609], [610], [611], [612], [613], [614], [615], [616], [617], [618], [619], [620], [621], [622], [623], [624], [625], [626], [627], [628], [629], [630], [631], [632], [633], [634], [635], [636], [637], [638], [639], [640], [641], [642], [643], [644], [645], [646], [647], [648], [649], [650], [651], [652], [653], [654], [655], [656], [657], [658], [659], [660], [661], [662], [663], [664], [665], [666], [667], [668], [669], [670], [671], [672], [673], [674], [675], [676], [677], [678], [679], [680], [681], [682], [683], [684], [685], [686], [687], [688], [689], [690], [691], [692], [693], [694], [695], [696], [697], [698], [699], [700], [701], [702], [703], [704], [705], [706], [707], [708], [709], [710], [711], [712], [713], [714], [715], [716], [717], [718], [719], [720], [721], [722], [723], [724], [725], [726], [727], [728], [729], [730], [731], [732], [733], [734], [735], [736], [737], [738], [739], [740], [741], [742], [743], [744], [745], [746], [747], [748], [749], [750], [751], [752], [753], [754], [755], [756], [757], [758], [759], [760], [761], [762], [763], [764], [765], [766], [767], [768], [769], [770], [771], [772], [773], [774], [775], [776], [777], [778], [779], [780], [781], [782], [783], [784], [785], [786], [787], [788], [789], [790], [791], [792], [793], [794], [795], [796], [797], [798], [799], [800], [801], [802], [803], [804], [805], [806], [807], [808], [809], [810], [811], [812], [813], [814], [815], [816], [817], [818], [819], [820], [821], [822], [823], [824], [825], [826], [827], [828], [829], [830], [831], [832], [833], [834], [835], [836], [837], [838], [839], [840], [841], [842], [843], [844], [845], [846], [847], [848], [849], [850], [851], [852], [853], [854], [855], [856], [857], [858], [859], [860], [861], [862], [863], [864], [865], [866], [867], [868], [869], [870], [871], [872], [873], [874], [875], [876], [877], [878], [879], [880], [881], [882], [883], [884], [885], [886], [887], [888], [889], [890], [891], [

fraction containing the receptor protein on the present a reaction vessel and incubate with the binding between the ligand and receptor protein. The reaction mixture is then separated by a method to prepare a receptor standard. Any buffer which does not interfere with the binding between the ligand and receptor protein can be used, for example, a phosphate buffer of pH 4 to 10, preferably pH 6 to 8, Tris-HCl buffer, or the like. In order to minimize non-specific binding, a protein extract such as CHAPS, Tween-20, Triton X-100, or the like, and various proteins such as bovine serum albumin, gelatin, etc. to the buffer. In addition, for inhibiting degradation of the receptor protein an agent by a protease, a protease inhibitor (e.g., EDTA, N-ethylmaleimide, phenylmethylsulfonyl fluoride, etc.) can also be added to the buffer. A given amount (5000 cpm-500,000 cpm) of the (peptide (enkeph-lysine), papain, etc. can also be added to the buffer. A given amount (5000 cpm-500,000 cpm) of the test compound labeled with [<sup>3</sup>H], [<sup>125</sup>I], [<sup>14</sup>C], [<sup>35</sup>S] or the like is added to 0.01 ml to 10 ml of the receptor suspension. For evaluating an amount of non-specific binding (NSB), a reaction tube, to which a large excess amount of the unlabeled test compound is added, is provided. The reaction is carried out at about 0°C to about 50°C, preferably about 4°C to about 37°C for about 20 minutes to about 24 hours, preferably about 30 minutes to about 3 hours. After completion of the reaction, the reaction mixture was filtered through, for example, glass fiber filter paper, washed with suitable amount of the same buffer and the radioactivity remaining in the glass fiber filter paper is counted with a liquid scintillation counter or a γ-counter. The test compound whose count (B-SB) obtained by subtracting the amount of non-specific binding (NSB) from the total binding amount (B) exceeds 0 cpm can be selected as the ligand to the receptor protein or as the seal of the present invention.

The above standard determination methods (d) and (e) can be carried out by measuring a set of intracellular  $\text{Ca}^{2+}$  activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $\text{Ca}^{2+}$  formation of intracellular cAMP, formation of intracellular cGMP, production of histoinl phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of  $\text{Ca}^{2+}$  ions, lowering of pH, etc.) mediated by the receptor of the present invention by a known method or a commercially available determination kit. Specifically, for the cells containing the receptor protein are cultivated in a well plate, and have cyclo-oxygenase. After addition of a test compound, the cells containing the receptor protein are cultured in a suitable buffer which does not have cyclo-oxygenase. The culture medium is replaced with a fresh medium or a suitable buffer of time, followed by extracting the cells or recovering the supernatant. The culture is incubated for a given period of time, followed by determining the product. Where an assay of the formation of a product formed by a method suitable for determination of the product. Where an assay of the formation of an index compound of a cell stimulating activity (e.g., arachidonic acid, etc.) is difficult due to a degradation enzyme contained in the cells, the assay can be carried out with addition of an inhibitor of the degradation enzyme. In addition, for cAMP production inhibitory activity or the like, the activity can be detected as the production inhibitory activity of cells whose basic production is increased with forskolin, etc.

The determination kit of a ligand which binds to the receptor protein or its salt of the present invention comprises as an essential component the receptor protein or the partial peptide of the present invention, cells containing the receptor protein of the present invention or a salt thereof, a cell membrane fraction of cells containing the receptor protein of the present invention, or the like.

Examples of the kid for ligand determination of the present invention include as follows

**(1) Reagent for ligand determination**

### Measurement buffer and washing buffer

A buffer obtained by addition of 0.05% of bovine serum albumin (Sigma) to distilled water (pH 7.4) (Bicco) can be prepared when it is used.

Standard of G-protein coupled receptor protein CHO cells expressing the receptor protein of the present invention are subjected to passage in an amount of  $1 \times 10^6$  cells/well in a 12-well plate and cultivated at 37°C for 2 days in 5% CO<sub>2</sub>-95% air to obtain a standard of receptor protein.

(c) Labeled test compound

A fast compound labeled with a commercial available [ $^3\text{H}$ ] ( $[^{3-}\text{H}]_1$ , [ $^{3-}\text{H}]_2$ , [ $^{3-}\text{H}]_3$ , [ $^{3-}\text{H}]_4$ , [ $^{3-}\text{H}]_5$ , etc., or unlabeled) compound is stored at  $4^\circ\text{C}$  or  $-20^\circ\text{C}$  and, when it is used, it is diluted.

An aqueous solution of the ladderized test compound is shown in Figure 1.

to 1  $\mu$ M with the measurement buffer. As for a water-insoluble or slightly water-soluble compound, the compound is dissolved in dimethylformamide, DMSO, methanol, etc.

(d) Non-labeled test compound

The same test compound as that of the labeled test compound is used to prepare a solution in the test cell at the same concentration.

**(2) Measurement**

(a) The receptor protein expressing CHO cells cultivated in a 12-well tissue culture plate is washed twice with ml of the measurement buffer and then 490  $\mu$ l of the buffer is added to the respective wells.

(b) 5.0 ml of the labeled test compound is added and reacted at room temperature for one hour

(c) The reaction mixture is removed and the wells are washed three times with 1 ml of the washing buffer. The labelled test compound bound to the cells is dissolved with 0.2 N NaOH-1% SDS and mixed with 4 ml of liquid scintillator A (Wako Pure Chemical Industries, Ltd.).

(d) Radioactivity is measured by a liquid scintillation counter (Beckman)

The ligand which can bind to the receptor protein of the present invention is, for example, that specifically present in brain, pituitary, pancreas, etc. Specific examples thereof include the above-exemplified Y, opiod, purine, vasopressin, bombesin, cannabinoid, cholecystic acid, glutamine, serotonin, melatonin, neuropeptide Y, opiod, putrine, vasopressin, oxytocin, PACAP, secretin, glucagon, calcitonin, dopaminomedullin, somatostatin, GHRP, CGF, ACTH, GRP, PTH, VIP, vasocadin, histamine and related polypeptides) dopamine, angiotensin II, substance P, galanin, neurokinin B, neurokinin A, neurokinin C, leukotriene, paracetamol, prostaglandin, thromboxane, adenosine, adrenalin,  $\alpha$ - $\beta$ -methionine (e.g., IL-1), prolactin, ghrelin, ghrelin, NPY-2, ENA-76, PAF, IP-10, GPCR-2, MCP-1, HC14, MCP-3, 1-309, MCP-18, RANTE, etc.; endothelin, angiotensin, histamine, neurotensin, TRH, pancreatic polypeptides or galinin.

(ii) Gene prophylactic and therapeutic drug of the G-protein coupled receptor protein deficiency disease

If a ligand to the receptor protein of the present invention is found out by the above method (1), DNA encoding the receptor protein of the present invention can be used as a drug for gene prophylaxis and therapy of diseases caused by deficiency of the G-protein coupled receptor protein deficiency depending upon a particular activity of the ligand.

For example, when there is a patient whose physiological activity of a ligand is scarcely expected because of a decrease in the receptor protein of the present invention, the amount of the receptor protein in the patient's body can be increased to exhibit the sufficient activity of the ligand, for example by (i) administering DNA encoding the receptor protein of the present invention to the patient to express it, or (ii) inserting DNA encoding the receptor protein of the present invention into subject cells to express it, followed by transplacental of the cells into the patient. Then, DNA encoding the receptor protein of the present invention is useful as a safe and low toxic drug for gene prophylaxis and therapy of disease caused by deficiency of the receptor protein of the present invention.

encapsulated in the receptor protein of the present invention (hereinafter referred to as the DNA or the present invention) alone or, after inserted into a suitable vector such as retroviral vector, adenoviral vector, adenoviral vector, etc., can be used as the above prophylactic and therapeutic drug according to a conventional method. For example, the DNA of the present invention can be used in the form of injectable preparations such as capsules, elixirs, microcapsules, etc. for oral administration, or in the form of injectable preparations such as cooling, capsules, elixirs, microcapsules, etc. for oral administration, or in the form of injectable preparations such as aseptic solutions or suspensions in water or other pharmaceutically acceptable solvents for parenteral administration. A pharmaceutical composition in a unit dosage form can be prepared by mixing the DNA of the present invention with pharmaceutically acceptable carriers, flavors, excipients, vehicles, preservatives, stabilizers, binders, etc. according to generally acceptable manner. The effective component is contained in the composition in such an amount that a dose in the intended desired range can be obtained.

Examples of additives to be mixed in tablets, capsules, etc. include binders such as gelatin, corn starch, tragacanth, croscarmellose, croscarmellose sodium, croscarmellose potassium, croscarmellose calcium, gum and gum arabic, excipients such as crystalline cellulose, swelling agents such as corn starch, gelatin and alginate, lubricants such as magnesium stearate, sweeteners such as sucrose, lactose and saccharin, flavors such as peppermint, aluminum oil and cherry, and the like. In case of the capsule dosage unit form, in addition to the above component, it can contain a liquid carrier such as fat or oil. An injectable aseptic composition can be prepared according to a conventional manner, for example, by dissolving the active component in a vehicle such as injectable water and a natural vegetable oil such as sesame oil, coconut oil, etc. Examples of the injectable aqueous solution include physiological saline, isotonic solutions containing glucose and other reagents (e.g., D-sorbitol, D-mannitol, sodium chloride, etc.) and suitable dissolution aids, for example, alcohols (e.g., ethanol), polyalkyls (e.g., polypropyl glycol, poly(ethylene glycol), nonionic surfactants (e.g., Polysorbate 80™, HCO-50) may be further added. As an oil solution, for example, sesame oil, soybean oil, etc. can be used and a dissolution aid such as benzyl benzoate can be added.

benzyl alcohol, etc. can be further added.

The above prophylactic and therapeutic drugs can further contain, for example, buffers (e.g., phosphate buffer, sodium acetate buffer), smoothing agents (e.g., benzalkonium chloride, procaine hydrochloride, etc.), stabilizers (e.g., human serum albumin, polyethylene glycol, etc.), preservatives (e.g., benzyl alcohol, phenol, etc.), antioxidants, and the like. The injectable preparation thus produced is normally filled in a suitable ampoule. Since the pharmaceutical composition thus obtained is safe and low toxic, it can be administered to a human being and another mammal (e.g., rat, rabbit, sheep, pig, cattle, cat, dog, monkey, etc.). Although the amount of the DNA of the present invention to be administered is varied according to particular subjects, organs to be treated, symptoms, routes of administration, etc., in general, for oral administration to an adult human being (as 60 kg body weight), the DNA is administered in an amount of about 0.1 mg/day to about 100 mg/day, preferably about 1.0 mg/day to about 50 mg/day, more preferably about 1.0 mg to about 20 mg. For parenteral administration to an adult human being (as 60 kg body weight), it is advantageous to administer the composition in the form of an injectable preparation in an amount of about 0.01 mg/day to about 30 mg/day, preferably about 0.1 mg/day to about 20 mg/day, more preferably about 0.1 mg/day to about 10 mg/day, though the single dosage is varied according to particular subjects, organs to be treated, symptoms, routes of administration, etc. As for other animals, the composition can be administered in the above amount with converting it into that for the body weight of 60 kg.

### (III) Gene diagnosing agent

The DNA encoding the receptor protein or the partial peptide of the present invention can be used for detecting an abnormality of DNA encoding the receptor protein or the partial peptide of the present invention (abnormal gene) in a human being or another mammal (e.g., rat, rabbit, sheep, pig, cattle, cat, dog, monkey, etc.). Therefore, the DNA encoding the receptor protein or the partial peptide of the present invention is useful as a gene diagnosing agent for detecting abnormality of the DNA.

### (IV) Determination method of ligand to the G-protein coupled receptor protein

The receptor protein, the partial peptide and their salts of the present invention have ligand binding properties and they can be used for determination of ligand concentration in the living body in a high sensitivity.

The method for determination of the present invention can be used, for example, in combination with a competitive method. That is, the ligand or concentration can be determined by bringing a specimen into contact with the receptor protein, the partial peptide or a salt thereof of the present invention. Specifically, the determination method can be carried out, for example, according to the methods described in Hiroshi Irie, Ed., "Radioimmunoassay", Kodan-sha, 1974 and "Second Series Radioimmunoassay", Kodan-sha, 1979 or their modifications.

### (V) Method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein

Compounds which alter binding of a ligand to the receptor protein or its salt (e.g., peptides, proteins, non-peptide compounds, synthetic compounds, fermentation products) can be screened efficiently by using the receptor protein, the partial peptide or their salts of the present invention, or by constructing an expression system of the recombinant receptor protein and using a receptor binding assay system utilizing the expression system. Examples of these compounds include compounds having cell stimulation activities (e.g., activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $Ca^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor protein of the present invention and compounds which do not have such activities (i.e., so-called antagonists to the receptor protein of the present invention).

That is, the present invention provides a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises comparing (i) binding of the ligand to the receptor protein or the partial peptide of the present invention or a salt thereof upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the ligand, and (ii) that upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the ligand and a test compound.

In the screening method of the present invention, an amount of the ligand bound to the receptor protein, the partial peptide or a salt thereof, a cell stimulation activity or the like is measured and compared upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the ligand, and (ii) that upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the ligand and a test compound.

More specifically, the screening method of the present invention includes:

- (a) a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to the receptor protein or the partial peptide of the present invention or a salt thereof upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the labeled ligand, and (ii) that upon bringing the receptor protein or the partial peptide of the present invention or a salt thereof into contact with the labeled ligand and a test compound;
  - (b) a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to cells containing the receptor protein or a membrane fraction of the cells upon bringing the labeled ligand into contact with the cells or the membrane fraction, and (ii) that upon bringing the labeled ligand and a test compound into contact with the cells or the membrane fraction;
  - (c) a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises labeling the ligand, and measuring and comparing (i) an amount of the labeled ligand bound to the receptor protein expressed on the cell membrane of a transformant containing DNA encoding the receptor protein of the present invention by culturing the transformant upon bringing the labeled ligand into contact with the expressed receptor protein, and (ii) that upon bringing the labeled ligand and a test compound into contact with the expressed receptor protein;
  - (d) a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises measuring and comparing (i) a cell stimulation activity (e.g., activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $Ca^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor upon bringing a compound which activates the receptor protein of the present invention (e.g., a ligand to the receptor protein of the present invention) into contact with cells containing the receptor protein of the present invention, and (ii) that upon bringing the compound which activates the receptor protein and a test compound into contact with the cells;
  - (e) a method for screening for compounds which alter binding of a ligand to the receptor protein or its salt of the present invention, or their salts which comprises measuring and comparing (i) a cell stimulation activity (e.g., activity to enhance or inhibit release of arachidonic acid, release of acetyl choline, release of intracellular  $Ca^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol phosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor upon bringing a compound which activates the receptor protein of the present invention (e.g., a ligand to the receptor of the present invention) into contact with the receptor protein expressed on the cell membrane of a transformant containing DNA encoding the receptor protein by culturing the transformant, and (ii) that upon bringing the compound which activates the receptor protein of the present invention and a test compound into contact with the receptor protein expressed on the cell membrane;
- Before the receptor protein of the present invention is available, for screening for a G-protein coupled receptor agonist or antagonist, first, it is necessary to obtain candidate compounds by using cells, tissues or a cell membrane fraction thereof containing the G-protein coupled receptor protein of rat, etc. (primary screening) and then to confirm whether or not the candidate compounds actually inhibit the binding of human G-protein coupled receptor protein and a ligand (secondary screening). When cells tissues or a cell membrane thereof are used as such, other receptor proteins are present, which make screening for an agonist or antagonist to the objective receptor protein difficult. However, for example, if the human receptor protein of the present invention is used, the primary screening is not required and efficient screening for compounds which inhibit the binding of a ligand to the G-protein coupled receptor protein can be carried out. In addition, whether the compound thus screened for is an agonist or an antagonist can be readily evaluated.
- The receptor protein used in the screening method of the present invention may be any protein in so far as it contains the above-described receptor protein or partial peptide of the present invention. A cell membrane fraction of a mammalian internal organ containing the receptor protein of the present invention is suitable. However, in particular, by using a recombinant is suitable for the screening.
- For producing the receptor protein of the present invention, the above-described expression process can be used. In particular, it is preferred to carry out expression of the DNA encoding the receptor protein in cells of mammals or insects. Normally, cDNA is used as a DNA fragment encoding the desired part of the protein, but the present invention is not necessarily limited to this. For example, a gene fragment or a synthetic DNA can also be used. For transferring



a DNA fragment encoding the receptor protein of the present invention into a host animal cell and expressing it efficiently, it is preferred to integrate the DNA fragment into the downstream from a polyprotein promoter of nuclear polyhedral virus (NPV) belonging to baculovirus whose host is insects, a promoter derived from SV40, retrovirus promoter, melibaculovirus promoter, human heat shock promoter, cytomegalovirus promoter, Spha promoter or the like. The quantitative and qualitative assay of the expressed receptor can be carried out by a per se known method. For example, the assay can be carried out according to the method described by Nambi, P. et al., J. Biol. Chem., 267, 19555-19559 (1992).

In the screening method of the present invention, the receptor protein, its partial peptide or their salts may be the receptor protein, its partial peptide or the salt as such which is purified according to a per se known method, or a material containing the receptor protein, its partial peptide or their salts such as cells containing the receptor protein or their membrane fractions.

In case of using the cells containing the receptor protein of the present invention, they may be immobilized with glutaraldehyde, formalin, etc., immobilization can be carried out by a per se known method.

The cells containing the receptor protein of the present invention are host cells expressing the receptor protein of the present invention. Preferred examples of the host cells include *E. coli*, *Bacillus subtilis*, yeast, insect cells, animal cells and the like.

The cell membrane fraction is a fraction containing many cell membranes obtained by disrupting cells and then treated by a per se known method. The disruption of cells can be carried out, for example, using Potter-Elvehjem homogenizer, Waring blender or Polytron (manufactured by Kinematica), ultrasonication, French press, etc. Fractionation of the cell membrane fraction can mainly be carried out by fractionation centrifugation, density gradient centrifugation or the like. For example, a disrupted cell suspension is centrifuged at a low rate (500 rpm-3000 rpm) for a short period of time (normally, about 1-10 minutes) and further the supernatant is centrifuged at a high rate (15000 rpm-30000 rpm) for about 30 minutes to about 2 hours to obtain a cell membrane fraction as a precipitate. This cell membrane fraction contains the expressed receptor protein and many membrane components such as phospholipids, membrane proteins and the like.

The amount of the receptor protein contained in the cells containing the receptor protein or the cell membrane fraction is preferably  $10^3$  to  $10^6$  molecules, more preferably  $10^5$  to  $10^7$  molecules per one cell. As the expression level is higher, a ligand binding activity (specific activity) becomes higher, which makes not only construction of a high sensitive screening system but also determination of a large number of samples in one lot possible.

For carrying out the above screening methods (a) to (c), a suitable receptor protein fraction and a labeled test compound are required. The receptor protein fraction is preferably a naturally occurring receptor protein, recombinant receptor protein fraction having the equivalent activity to that of the naturally occurring receptor protein. The equivalent activity used herein means equivalent ligand binding activity, signal information transmission activity or the like.

The labeled ligand includes labeled ligands and labeled ligand analog compounds. For example, the ligand labeled with [ $^3$ H], [ $^{125}$ I], [ $^{14}$ C], [ $^{35}$ S], etc., can be used.

Specifically, for carrying out the screening method of the present invention, the cells or cell membrane fraction containing the receptor protein of the present invention are suspended in a buffer suitable for the screening to prepare a receptor protein standard. Any buffer which does not interfere with the binding between the ligand and the receptor protein can be used, for example, a phosphate buffer of pH 4 to 10, preferably pH 6 to 8, Tris-HCl buffer or the like. In order to minimize non-specific binding, a surfactant such as CHAPS, Tween-20™ (Kao-Alfa), digitonin, deoxycholate, etc., and various proteins such as bovine serum albumin, gelatin, etc., to the buffer. In addition, for inhibiting degradation of the receptor protein and the ligand by a protease, a protease inhibitor such as PMSF, leupeptin, E-64 (manufactured by Peptide Kenkyu-sho), pepstatin, etc., can also be added to the buffer. At the same time,  $10^{-4}$  M of cpm-500000 (cpm) is added to 0.01 ml to 10 ml of the receptor suspension and, at the same time,  $10^{-4}$  M of a test compound is present in the receptor suspension. For evaluating an amount of non-specific binding (NSB), a reaction tube to which a large excess amount of the unlabeled ligand is added is provided. The reaction is carried out about 0°C to about 50°C, preferably about 4°C to about 37°C for about 20 minutes to about 24 hours, preferably about 30 minutes to about 3 hours. After completion of the reaction, the reaction mixture was filtered through, for example, a glass fiber filter paper, washed with a suitable amount of the same buffer and the radioactivity remaining in the glass fiber filter paper is counted with a liquid scintillation counter or a  $\gamma$ -counter. The test compound whose specific binding count (B-NSB) is 50% or lower can be selected as a candidate compound capable of inhibiting antagonist, when taking the count (B<sub>0</sub>-NSB) obtained by subtracting the count of non-specific binding (NSB) from the count (B<sub>0</sub>) obtained without any antagonistic material as 100%.

The above screening methods (d) and (e) can be carried out by measuring a cell stimulation activity (e.g., activity to enhance or inhibit release of arachidonic acid, release of acetylcholine, release of intracellular  $Ca^{2+}$ , formation of intracellular cAMP, formation of intracellular cGMP, production of inositol triphosphate, change of cell membrane potential, phosphorylation of intracellular protein, activation of c-fos, lowering of pH, etc.) mediated by the receptor protein of the

present invention by a known method or a commercially available determination kit. Specifically, first, the cells containing the receptor protein are cultivated in a multi-well plate, etc. For carrying out screening, the culture medium is replaced with a fresh medium or a suitable buffer which does not have cytotoxicity. After addition of a test compound, the culture is incubated for a given period of time, followed by extracting the cells or recovering the supernatant to determine a product formed by a method suitable for determination of the product. Where an assay of the formation of an index compound of a cell stimulating activity (e.g., arachidonic acid, etc.) is difficult due to a degradation enzyme contained in the cells, the assay can be carried out with addition of an inhibitor of the degradation enzyme. In addition, as for cAMP production inhibitory activity or the like, the activity can be detected as the production inhibitory activity of cells whose basic production has been increased with forskolin, etc.

For carrying out the screening by measurement of a cell stimulating activity, cells expressing a suitable receptor protein are required. As cells expressing the receptor protein of the present invention, for example, naturally occurring cell strains containing the receptor protein of the present invention or the above-described recombinants expressing the receptor protein are preferred.

Examples of the test compounds include peptides, proteins, non-peptide compounds, synthetic compounds, cell extracts, vegetable extracts, animal tissue extracts and the like and these compounds may be novel compounds or known compounds.

The kit for screening for a compound which alters binding of a ligand to the receptor protein or its salt of the present invention comprises as an essential component the receptor protein or the partial peptide of the present invention or a salt thereof, cells containing the receptor protein of the present invention, a cell membrane fraction of cells containing the receptor protein of the present invention, or the like.

Examples of the screening kit of the present invention include as follows.

#### (1) Reagent for screening

##### (a) Measurement buffer and washing buffer

A buffer obtained by addition of 0.05% of bovine serum albumin (Sigma) to Hank's balanced salt solution (Gibco). The buffer is sterilized by filtration through a filter of 0.45  $\mu$ m pore diameter and then stored at 4°C or is can be prepared when it is used.

##### (b) Standard of G-protein coupled receptor protein

CHO cells expressing the receptor protein of the present invention are subjected to passage in an amount of  $5 \times 10^5$  cells/well in a 12-well plate and cultured at 37°C for 2 days in 5% CO<sub>2</sub>-95% air to obtain a standard of the receptor protein.

##### (c) Labeled ligand

A ligand labeled with a commercially available [ $^3$ H], [ $^{125}$ I], [ $^{14}$ C], [ $^{35}$ S], etc., or another suitable label. An aqueous solution of the labeled test compound is stored at 4°C or -20°C and, when it is used, it is diluted to 1  $\mu$ M with the measurement buffer.

##### (d) Ligand standard solution

A ligand is dissolved in PBS containing 0.1% bovine serum albumin (Sigma) at the final concentration of 1 mM and stored at -20°C.

#### (2) Measurement

(a) The receptor protein expressing CHO cells cultivated in a 12-well tissue culture plate is washed twice with 1 ml of the measurement buffer and then 450  $\mu$ l of the same buffer is added to the respective wells.

(b) 5  $\mu$ l of a  $10^{-3}$  to  $10^{-10}$  M solution of a test compound is added and then 5  $\mu$ l of the labeled ligand is added. They are reacted at room temperature for one hour. In order to evaluate a non-specific binding amount, 5  $\mu$ l  $10^{-3}$  M ligand is added.

(c) The reaction mixture is removed and the wells are washed three times with 1 ml of the washing buffer. The labeled ligand bound to the cells is dissolved with 0.2 N NaOH-1% SDS and mixed with 4 ml of liquid scintillator A (Mako Pure Chemical Industries, Ltd.).

(d) Radioactivity is measured by a liquid scintillation counter (Beckman). Percent Maximum Binding (PMB) is calculated by the equation [1]:

$$PMB = (B - NSB) / (B_0 - NSB) \times 100$$

wherein PMB is Percent Maximum Binding, B is a value obtained with addition of a specimen, NSB is a value of non-specific binding, and B<sub>0</sub> is a maximum binding value.

The compound or its salt obtained by using the screening method of the present invention alters binding of a ligand to the receptor protein or its salt of the present invention. Specifically, it is a compound or a salt thereof which binds to the receptor protein of the present invention and exhibits cell stimulating activity mediated by the receptor protein (i.e., so-called an agonist to the receptor protein of the present invention) or which binds to the receptor protein of the present invention but does not exhibit the cell stimulating activity (i.e., so-called an antagonist of the present invention).

Examples of the compounds include peptides, proteins, non-peptide compounds, synthetic compounds, fermented products and the like and they may be novel compounds or known compounds.

Since the agonist to the receptor protein of the present invention has the same physiological activity as that of a ligand to the receptor protein, it is useful as an active component of a safe and low toxic pharmaceutical composition having the ligand activity.

On the other hand, since the antagonist to the receptor protein of the present invention can inhibit the physiological activity of a ligand to the receptor protein, it is useful as an active component of a safe and low toxic pharmaceutical composition for inhibiting the ligand activity.

If the compound or its salt obtained by the screening method or the screening kit of the present invention is used for a pharmaceutical composition as described above, any conventional manner can be employed. For example, the compound or its salt of the present invention can be used in the form of tablets, if necessary, providing sugar coating, capsules, elixirs, microcapsules, etc. for oral administration, or in the form of injectable preparations such as aseptic solutions or suspensions in water or other pharmaceutically acceptable solutions for parenteral administration. A pharmaceutical composition in a unit dosage form can be prepared by mixing the compound or its salt of the present invention with, for example, one or more pharmaceutically acceptable carriers, flavors, excipients, vehicles, preservatives, stabilizers, binders, etc. according to generally acceptable manner. The effective component is contained in the composition in such an amount that a dose in the intended desired range can be obtained.

Examples of additives to be mixed in tablets, capsules, etc. include binders such as gelatin, corn starch, tragacanth gum and gum arabic, excipients such as crystalline cellulose, swelling agents such as corn starch, gelatin and alginate acid, lubricants such as magnesium stearate, sweeteners such as sucrose, lactose and saccharin, flavors such as peppermint, alkamono oil and cherry, and the like. In case of a capsule dosage unit form, in addition to the above components, it can contain a liquid carrier such as fat or oil. An injectable aseptic composition can be prepared according to a conventional manner, for example, by dissolving or suspending the active component in a vehicle such as injectable water and a natural vegetable oil such as sesame oil, coconut oil, etc. Examples of injectable aqueous solution include physiological saline, isotonic solutions containing glucose and other adjuvants (e.g., D-sorbitol, D-mannitol, sodium chloride, etc.) and suitable dissolution aids, for example, alcohols (e.g., ethanol), polyalcohols (e.g., propylene glycol, polyethylene glycol), nonionic surfactants (e.g., polyoxyethylene glycol, HCO-50) may be further added. As an oily solution, for example, sesame oil, soybean oil, etc. can be used and a dissolution aid such as benzyl benzoate or benzyl alcohol, etc. can be further added. The composition can further contain, for example, buffers (e.g., phosphate buffer, sodium acetate buffer), smoothing agents (e.g., benzalkonium chloride, procaine hydrochloride, etc.), stabilizers (e.g., human serum albumin, polyethylene glycol, etc.), preservatives (e.g., benzyl alcohol, phenol, etc.), antioxidants, and the like. The injectable preparation thus produced is normally filled in an appropriate ampoule.

Since the pharmaceutical composition thus obtained is safe and low toxic, it can be administered to a human being and another mammal (e.g., rat, rabbit, sheep, pig, cattle, cat, dog, monkey, etc.).

Although the amount of the compound or its salt of the present invention to be administered is varied according to particular subjects, internal organs to be treated, symptoms, routes of administration, etc. in general, for oral administration to an adult human being (as 60 kg body weight), the DNA is administered in an amount of about 0.1 mg/day to about 100 mg/day, preferably about 1.0 mg/day to about 50 mg/day, more preferably about 1.0 mg to about 20 mg. For parenteral administration to an adult human being (as 60 kg body weight), it is advantageous to administer the composition in the form of an injectable preparation in an amount of about 0.01 mg/day to about 30 mg/day, preferably about 0.1 mg/day to about 20 mg/day, more preferably about 0.1 mg/day to about 10 mg/day, though the single dosage is varied according to particular subjects, internal organs to be treated, symptoms, routes of administration, etc. As for other animals, the composition can be administered in the above amount with converting it into that for the body weight of 60 kg.

(VI) Preparation of antibody or antiserum against the G-protein coupled receptor protein, the partial peptide or salts thereof

An antibody (e.g., monoclonal antibody, polyclonal antibody) or antiserum against the receptor protein, the partial peptide or their salts of the present invention can be prepared by using the receptor protein or the partial peptide of the present invention or a salt thereof as the antigen according to a conventional antibody or antiserum preparation process. For example, a monoclonal antibody can be prepared as follows.

# Preparation of monoclonal antibody

## (a) Preparation of monoclonal antibody producer cells

The receptor protein or the partial peptide of the present invention or a salt thereof (hereinafter sometimes abbreviated to the receptor protein, etc.) as such or together with a suitable carrier or diluent is administered to a site of a mammal which permits the antibody production. For enhancing the antibody production capability, complete Freund's adjuvant or incomplete Freund's adjuvant may be administered. Normally, the receptor, etc. is administered once every 3 weeks to 6 weeks, in total, about 2 to about 10 times. The mammal to be used include monkey, rabbit, dog, guinea pig, mouse, rat, sheep, goat, chicken and the like and mouse or rat is preferred.

For preparing monoclonal antibody producer cells, an individual whose antibody titer has been confirmed is selected from warm blood animals immunized with the antigen, for example, rat and, 2 days to 5 days after the final immunization, its spleen or lymph node is collected and antibody producer cells contained therein are fused with myeloma cells to prepare the desired monoclonal antibody producer hybridoma. Measurement of the antibody titer in an antiserum can be carried out, for example, by reacting the labeled receptor protein, etc. as described hereinafter and an antiserum and then measuring the activity of the labeling agent bound to the antibody. The cell fusion can be carried out according to a known method, for example, the method described by Kohler and Milstein, *Nature*, 255, 495 (1975). As a fusion promoter, for example, polyethylene glycol (PEG) or Sendai virus (HVJ), preferably PEG can be used.

Examples of myeloma cells include NS-1, P3U1, SP20, AP-1 and the like and P3U1 is preferred. The proportion of the number of antibody producer cells (spleen cells) and the number of myeloma cells to be used is preferably about 1:1 to about 20:1 and PEG (preferably PEG 1000-PEG 6000) is added in concentration of about 10% to about 80%. Cell fusion can be carried out efficiently by incubating a mixture of both cells at about 20°C to about 40°C, preferably about 30°C to about 37°C for about 1 minute to about 10 minutes.

Various methods can be used for screening for a hybridoma producing the antibody against the receptor protein, etc. For example, there may be mentioned a method wherein a supernatant of the hybridoma is added to a solid phase (e.g., microplate) to which the receptor protein antibody is adsorbed directly or together with a carrier and then an anti-immunoglobulin antibody (if cells of a mouse is used in cell fusion, anti-mouse immunoglobulin antibody is used) of Protein A labeled with a radioactive substance or an enzyme is added to detect the monoclonal antibody against the receptor protein, etc. bound to the solid phase, and a method wherein a supernatant of the hybridoma is added to a solid phase to which an anti-immunoglobulin antibody or Protein A is adsorbed and then the receptor protein, etc., labeled with a radioactive substance or an enzyme is added to detect the monoclonal antibody against the receptor protein, etc. bound to the solid phase.

Selection of the monoclonal antibody can be carried out according to a per se known method or its modification. Normally, a medium for animal cells to which HAT (hypoxanthine, aminopterin, thymidine) are added is employed. Any selection and growth medium can be employed in so far as the hybridoma can grow. For example, RPMI 1640 medium containing 1% to 20%, preferably 10% to 20% fetal bovine serum, GIT medium containing 1% to 10% fetal bovine serum (Wako Pure Chemical Industries, Ltd.), a serum free medium for cultivation of a hybridoma (SFM-101, Nissui Seiyaku) and the like can be used. Normally, the cultivation is carried out at 20°C to 40°C, preferably 37°C for about 5 days to about 3 weeks, preferably 1 week to 2 weeks under about 5% CO<sub>2</sub> gas. The antibody titer of the supernatant of a hybridoma culture can be measured according to the same manner as described above with respect to the antibody titer of the anti-G-protein coupled receptor in the antiserum.

## (b) Purification of monoclonal antibody

Separation and purification of a monoclonal antibody against the receptor protein, etc. (hereinafter sometimes referred to as the anti-receptor protein monoclonal antibody) can be carried out according to the same manner as those of conventional polyclonal antibodies such as separation and purification of immunoglobulins, for example, salting-out, alcoholic precipitation, isoelectric point precipitation, electrophoresis, adsorption and desorption with ion exchangers (e.g., DEAE), ultracentrifugation, gel filtration, or a specific purification method wherein only an antibody is collected with an active adsorbent such as an antigen-binding solid phase, Protein A or Protein G and dissociating the binding to obtain the antibody.

The antibody against the receptor protein, etc. of the present invention prepared by the above (a) and (b) can specifically recognize the receptor protein, etc. of the present invention and therefore it can be used in a quantitative determination of the receptor protein, etc. of the present invention in a specimen, particularly, a quantitative determination by a sandwich immunoassay. That is, the present invention also provides:

(a) a method for determination of the receptor protein or its salt of the present invention in a specimen fluid which comprises reacting an antibody against the receptor protein or the partial peptide of the present invention, the

specimen fluid and the labeled receptor protein or the labeled partial peptide competitively and measuring the proportion of the labeled receptor protein or the labeled partial peptide bound to the antibody; and  
(b) a method for determination of the receptor protein or its salt of the present invention in a specimen fluid which comprises reacting the specimen fluid, an antibody insolubilized on a carrier and a labeled antibody simultaneously or continuously, and then measuring the activity of the labeling agent on the insolubilized carrier, one antibody being that recognizing the N-terminal of the receptor protein of the present invention and the other antibody being that reacting with the C-terminal of the receptor protein of the present invention.

In addition to the determination of the receptor protein or its salt of the present invention, the anti-receptor protein monoclonal antibody recognizing the receptor protein of the present invention, etc. can be used for detection by histological status and the like. For these purposes, the antibody molecular as such can be used or Fab<sub>2</sub>, Fab' or Fab fragments of the antibody molecule can also be used. A method for determination using an antibody against the receptor protein, etc. of the present invention is not specifically limited and any determination method can be used in so far as an amount of an antigen, antibody or antibody-antigen corresponding to an amount of an antigen in a fluid to be determined can be detected by a chemical or physical means and calculated based on a calibration curve prepared by using standard solutions containing known amounts of the antigen. For example, nephelometry, competitive method, immunometric method and sandwich method are suitably employed. In particular, in view of sensitivity, specificity and immunometric method as described hereinafter is preferred.

As a labeling agent used in a determination method using a labeled reagent, radioisotopes, enzymes, fluorescent materials, luminous materials and the like can be used. Examples of radioisotopes include <sup>125</sup>I, <sup>131</sup>I, <sup>125</sup>Na, <sup>125</sup>Na, <sup>125</sup>Na and the like. As the above enzymes, that having good stability and high specific activities is preferred and, for example, there are  $\beta$ -galactosidase,  $\beta$ -glucosidase, alkaline phosphatase, peroxidase, male dehydrogenase and the like. As the fluorescent materials, for example, there are fluorescein, fluorescein isothiocyanate and the like. As the luminous materials, there are luminol, luminol derivatives, luciferin, luciferin and the like. In addition, biotin-avidin system can be used for binding of an antibody or antigen to a labeling agent.

For insolubilization of an antigen or antibody, physical adsorption can be used or, normally, a method using a chemical bond for insolubilizing or immobilizing a protein or an enzyme can be used. Examples of the carrier include insoluble polysaccharides such as agarose, dextran, cellulose and the like, synthetic resins such as polystyrene, polyacrylamide, silicone and the like, glass and the like.

In a sandwich method, a specimen fluid to be tested is reacted with an insolubilized anti-receptor protein antibody (primary reaction) and further reacting a labeled anti-receptor protein antibody (secondary reaction), followed by measuring the labeling agent on the insoluble carrier to determine the amount of the receptor protein of the present invention in the specimen. The order of the primary and secondary reactions can be reversed and they can be carried out simultaneously or separately at different times. The above-described labeling agent and insolubilization can be applied to this method. In addition, in an immunoassay by a sandwich method, an antibody to be used as the solid phase antibody or labeled antibody is not necessary one kind of antibodies and, in order to improve measuring sensitivity, etc., a mixture of two or more kinds of antibodies can be used.

In the method for determination of the receptor protein, etc. by the sandwich method of the present invention, preferably, the antibodies against the receptor protein, etc. used in the primary and secondary reactions are those having different binding sites for the receptor protein. For example, when the antibody used in the secondary reaction that recognizing the C-terminal region of the receptor protein, the antibodies used in the primary reaction is that recognizing an region other than the C-terminal region, for example, the N-terminal region.

The antibody against the receptor protein, etc. of the present invention can also be used for a measuring system other than a sandwich method, for example, a competitive method, immunometric method, nephelometry and the like. In a competitive method, an antigen in a specimen fluid and a labeled antigen are reacted with the antibody competitively and, after separation of the unreacted labeled antigen (F) from the labeled antigen bound to the antibody (B), measuring the amount of the labeling agent of either B or F to determine the amount of the antigen in the specimen fluid. In this reaction, both liquid phase method and solid phase method can be employed. In the liquid phase method, a soluble antibody is used as the antibody and B/F separation can be carried out by using polyethylene glycol, a second antibody, against the above antibody. In the solid phase method, an immobilized solid phase antibody is used as the first antibody, or a soluble antibody is used as the first antibody and an immobilized solid phase antibody is used as the second antibody.

In the immunometric method, an antigen in a specimen fluid to be tested and an immobilized solid phase antigen are reacted with a given amount of a labeled antibody, competitively and then the solid phase is separated from the liquid phase. Alternatively, an antigen in a specimen fluid to be tested is reacted with an excess amount of a labeled antibody and an immobilized solid phase antigen is added to permit the unreacted labeled antibody to bind to the solid phase, followed by the separation of the solid phase from the liquid phase. Then, the amount of the labeling agent of either phase is measured to determine the amount of the antigen in the specimen fluid.

In nephelometry, an antigen-antibody reaction is carried out in a gel or solution and the amount of an insoluble precipitate formed is measured. Even when the amount of an antigen in a specimen fluid to be tested is small and the amount of a precipitate formed is small, laser nephelometry wherein diffusion of laser is utilized can be suitably employed.

When employing these immunoassay methods in the determination method of the present invention, to set any special conditions, procedures and the like is not required. That is, the determination system of the receptor protein or its salt of the present invention can be constructed based on conventional conditions and procedures in respective methods together with conventional artian's technical consideration. As for details of these general technical means, reference can be made to various reviews, texts and the like, for example, Hiroshi Ito, Ed. Radioimmunoassay, Kodansha (1974); Hiroshi Ito, Ed. Second Series, Radioimmunoassay, Kodansha (1979); Eizi Ishikawa et al., Ed. Enzyme immunoassay, Igaku-shoin (1978); Eizi Ishikawa et al., Ed. Second Series, Enzyme immunoassay, Igaku-shoin (1982); Eizi Ishikawa et al., Third Series, Enzyme immunoassay, Igaku-shoin (1987). Method in Enzymology, Vol. 70, Immunochemical Techniques (Part C), ibid., Vol. 84, Immunochemical Techniques (Part D, Selected Immunoassays), ibid., Vol. 92, Immunochemical Techniques (Part E, Monoclonal Antibodies and General Immunoassay Methods), ibid., Vol. 121, Immunochemical Techniques (Part I, Hybridoma Technology and Monoclonal Antibodies and the like).

As described hereinafter, the receptor protein or its salt of the present invention can be determined at high sensitivity by using an antibody against the receptor protein, etc. of the present invention.

In the specification and drawings, the abbreviations of bases, amino acids and the like are those according to IUPAC-IUB Commission on Biochemical Nomenclature or those conventionally used in the art. The examples are as follows. When the amino acid has an optical isomer, the amino acid is L-isomer unless otherwise stated.

20	DNA:	deoxyribonucleic acid
	cDNA:	complementary deoxyribonucleic acid
25	A:	adenine
	T:	thymine
	G:	guanine
	C:	cytosine
30	mRNA:	messenger ribonucleic acid
	dATP:	deoxyadenosine triphosphate
	dTTP:	deoxythymine triphosphate
	dGTP:	deoxyguanosine triphosphate
35	dCTP:	deoxycytidine triphosphate
	ATP:	adenosine triphosphate
	EDTA:	ethylenediaminetetraacetic acid
	SDS:	sodium dodecylsulfate
	EIA:	enzyme immunoassay
40	Gly:	glycine
	Ala:	alanine
	Val:	valine
	Leu:	leucine
	Ile:	isoleucine
45	Ser:	serine
	Thr:	threonine
	Cys:	cysteine
	Met:	methionine
	Glu:	glutamic acid
	Asp:	aspartic acid
50	Lys:	lysine
	Arg:	arginine
	His:	histidine
	Phe:	phenylalanine
55	Tyr:	tyrosine
	Trp:	tryptophan
	Pro:	proline
	Asn:	asparagine
	Gln:	glutamine

pGlu : pyroglutamic acid  
 Me : methyl group  
 Et : ethyl group  
 Bu : butyl group  
 Ph : phenyl group  
 TC : thiazolidin-4(R)-carboxamide group

The sequences in the Sequence Listing of the present specification are as follows.

10 SEQ ID NO: 1

This represents an amino acid sequence of the G-protein coupled receptor protein derived from a human brain.

15 SEQ ID NO: 2

This represents an amino acid sequence of the G-protein coupled receptor protein derived from a human brain, which is a variant of SEQ ID NO: 1 having additional 61 amino acids at the N-terminal thereof.

20 SEQ ID NO: 3

This represents a nucleotide sequence of DNA encoding the G-protein coupled receptor protein derived from a human brain having the amino acid sequence represented by SEQ ID NO: 1.

25 SEQ ID NO: 4

This represents a nucleotide sequence of DNA encoding the G-protein coupled receptor protein derived from a human brain having the amino acid sequence represented by SEQ ID NO: 2, which is a variant of SEQ ID NO: 3 having additional 183 bases at the 5'-terminal thereof.

30 SEQ ID NO: 5

A nucleotide sequence of EST which has been registered with a data base (NCBI abEST) under the accession number of T08099.

35 SEQ ID NO: 6

A nucleotide sequence of EST which has been registered with a data base (NCBI abEST) under the accession number of T27053.

40 SEQ ID NO: 7

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

45 SEQ ID NO: 8

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

50 SEQ ID NO: 9

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

55 SEQ ID NO: 10

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

SEQ ID NO: 11

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

5 SEQ ID NO: 12

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

10 SEQ ID NO: 13

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

15 SEQ ID NO: 14

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

20 SEQ ID NO: 15

This represents a nucleotide sequence of a synthetic DNA used in screening for cDNA encoding the G-protein coupled receptor protein of the present invention.

25 The transformant Escherichia coli HB101/pHEBF2 obtained in Example 1 hereinafter has been deposited with National Institute of Bioscience and Human-Technology (NIBH), Agency of Industrial Science & Technology Ministry of International Trade & Industry (1-3, Higashi 1-chome, Tsukubashi Ibaraki, 305 Japan) according to the Budapest Treaty under the accession number of FERM BP-5724 since October 25, 1996 and also deposited with Institute for Fermentation Osaka (IFO, 17-85, Jusoh-honmachi 2-chome Yodogawa-ku, Osaka, 532 Japan) under accession number of IFO 16044 since October 21, 1996.

30 The following examples further illustrate the present invention in detail but are not to be construed to limit the scope thereof.

#### Example 1

Obtaining cDNA of entire translated region of receptor protein from human poly(A)<sup>+</sup>RNA and sequence analysis of its nucleotide sequence

(1) Obtaining cDNA of translated region about C-terminal of the receptor protein from human fetal brain poly(A)<sup>+</sup>RNA by 3' RACE (Rapid Amplification of cDNA End) method and sequence analysis of its nucleotide sequence in order to obtain cDNA of the translated region about the C-terminal of the receptor protein encoded by the known nucleotide sequences, accession numbers of T08099 (SEQ ID NO: 5) and T27053 (SEQ ID NO: 6). 3' RACE method was carried out by using human fetal brain poly(A)<sup>+</sup>RNA as a PCR template.

First, the following two primers were synthesized based on the known nucleotide sequences.

B1: (SEQ ID NO: 7)

5' -AAGTTGGCTGTCTCATCTGGTGGGCTC-3'

B2: (SEQ ID NO: 8)

5' -TGAGCTCTCTGCTGTGGCAGTGGCAG-3'

Then, a PCR template of 3' RACE method was prepared from 1 µg human fetal poly(A)<sup>+</sup>RNA (Clontech) by using 3' RACE kit (Gibco BRL). The first PCR was carried out using the primer attached to 3' RACE kit and B1 primer. The conditions were 30 seconds at 95°C, 60 seconds at 65°C and 180 seconds at 72°C, for 35 cycles and



Ex. Taq (Takara Shuzo) was used as the DNA polymerase. The second PCR reaction was carried out by using 1 µl of the first PCR mixture as the template under the same conditions for 35 cycles except that B2 primer was used instead of B1 primer. After electrophoresis, a band formed of 1.5 kb was recovered, subcloned by using a TA cloning kit (Invitrogen) and transferred into *E. coli* JM109. As a result of sequence analysis, it was found that the amplified band had the C-terminal region of the above-described known nucleotide sequences.

(2) Obtaining cDNA of translated region about N-terminal of the receptor protein from human fetal brain poly (A)<sup>+</sup>RNA by 5' RACE method (invention method) and sequence analysis of its nucleotide sequence. In order to obtain cDNA of the translated region about the N-terminal of the receptor protein encoded by the above-described known nucleotide sequences, 5' RACE method was carried out by using human fetal brain poly (A)<sup>+</sup> RNA as a PCR template.

First, the following two primers were synthesized based on the above-described known nucleotide sequences.

B8: (SEQ ID NO: 9)

5'-CATCGGGCGCTCTGTGTAGTATCATC-3'

B9: (SEQ ID NO: 10)

5'-GAGAGGATGGGCGAGCGAGAGTAGAG-3'

Then, a PCR template of 5' RACE method was prepared from 1 µg human fetal poly(A)<sup>+</sup>RNA (Clontech) according to the manual of 5' RACE kit (Clontech). The first PCR was carried out using the primer attached to 5' RACE kit and B9 primer. The conditions were 10 seconds at 98°C and 180 seconds at 72°C, for 5 cycles; 10 seconds at 98°C and 180 seconds at 70°C, for 5 cycles; and 10 seconds at 98°C and 180 seconds at 68°C, for 35 cycles. Ex Taq (Takara Shuzo) was used as the DNA polymerase. The second PCR reaction was carried out by using 1 µl of 50-fold dilution of the first PCR mixture as the template under the same conditions except that B8 primer was used instead of B9 primer. After electrophoresis, a band formed of about 1 kb was recovered and subcloned by using a TA cloning kit (Invitrogen), and sequence analysis was carried out according to the same manner as in the above (1). The presence of the 7-transmembrane receptor protein comprising 481 amino acids (SEQ ID NO: 1) encoded by the nucleotide sequence of from the 625th base (ATG, Met) to the 2067th base (TGC, Cys) as shown by Fig. 1 has been confirmed based on the results of the above (1) and (2). Fig. 2 shows the result of hydrophobic plotting of this amino acid sequence.

(3) The N-terminal side of the above receptor protein was further examined and 5' RACE method at the 5' side was further carried out to determine a transcription initiation codon.

The following two primers were synthesized based on the translation region of the receptor protein obtained in the above (2).

B11: (SEQ ID NO: 11)

5'-ATGAGGGCGACGCGACGAGAGAAAGC-3'

B12: (SEQ ID NO: 12)

5'-ATGACAAATAGGAGGCGAGAGAAAGAG-3'

According to the same manner as in the above (2), a PCR template was prepared by using the above human fetal brain poly(A)<sup>+</sup>RNA (Clontech) or human cerebellum poly(A)<sup>+</sup>RNA (Nippon gene). Then, according to the same manner as described in the above (2), twice PCR were carried out by using the primer B11 instead of the primer B9 and the primer B12 instead of the primer B8, respectively to amplify the templates derived from the poly(A)<sup>+</sup>RNAs of the above internal organs. After electrophoresis of the reaction product, a band formed was recovered, subcloned by TA cloning and then subjected to sequence analysis.

As a result, the sequence obtained by combining the above (1) to (3) had the nucleotide sequence (SEQ ID NO:

4) of from the 442nd base (ATC, Met) to the 2067th base (TGC, Cys) as shown in Fig. 3 and was confirmed that it encoded the 7-transmembrane receptor protein comprising 542 amino acids (SEQ ID NO: 2). The result of the hydrophobic plotting of this sequence is shown in Fig. 4.

The following primers were synthesized based on this sequence.

HEF: (SEQ ID NO: 13)

5'-GTGACGAGATGTGTAGAGGCGAGCAAGAGTGC-3'

HER-1: (SEQ ID NO: 14)

5'-TACTGGGGCCTCAGCAGAGGTGTGCCAG-3'

PCR was carried out by using these two primers to amplify the coding region for the receptor protein from human fetal brain cDNA library. After subcloning in *E. coli*, a clone without any PCR error was selected and transformed into *E. coli* HB101 to obtain the transformant, *E. coli* HB101/pHEF-2. The DNA contained in the plasmid pHEF-2 has the nucleotide sequence represented by SEQ ID NO: 4 (Fig. 3) and the nucleotide sequence represented by SEQ ID NO: 3 is contained therein.

When PCR was carried out by using the above two primers under the conditions of 30 seconds at 95°C and 90 seconds at 68°C to amplify the coding region, the cDNA of the receptor protein as shown by Fig. 3 was obtained from fetal brain.

In order to obtain the cDNA of the receptor protein from adult brain, HEF-2 primer was synthesized instead of HEF primer.

HEF-2: (SEQ ID NO: 15)

5'-GTCGACTGGCTGTCTCTGCTCATTCAGGCAT-3'

When adult brain poly(A)<sup>+</sup>RNA was amplified by using the primers HEF-2 and HEF-1, the cDNA encoding the receptor protein shown in Fig. 1 was obtained. The N-terminal of this receptor protein was shorter than that of the receptor protein shown in Fig. 3 by 61 amino acids.

The receptor protein shown in Fig. 1 has a consensus sequence called as Coxsack sequence which indicates initiation of translation just before the translation initiation codon and a defined signal sequence in the N-terminal region. In view of these results, it has been found that the receptor protein shown by Fig. 1 is predominantly expressed in adult brain. On the other hand, cDNA encoding the receptor protein shown by Fig. 3 is obtained from fetal brain and the presence of the long chain receptor protein shown by Fig. 3 has also been confirmed.

#### Example 2

Confirmation of expression specifically in various tissues

Northern blot was carried out by using cDNA encoding the receptor protein of the present invention contained in the plasmid pHEF-2 obtained in Example 1 as a probe.

The cDNA was labeled with Amersham's multiprime kit and [32P]dCTP according to the manual of the kit. Hybridization was carried out by using human MTN BclI (Clontech) and according to the manual attached thereto. Exposure of the filter to light was made as -80°C for one week. As shown in Fig. 5, it has been found that this receptor protein mRNA is specifically expressed in brain.

As described hereinabove, the G-protein coupled receptor protein, its partial peptide or their salts and DNAs encoding them of the present invention can be used (a) for determination of ligands, (b) for obtaining antibodies and antisera, (c) for construction of recombinant receptor protein expression systems, (d) for development of receptor binding assay systems and screening for drug candidate compounds by using the expression systems, (e) for practice of drug design based on comparison of structurally analogous ligand receptors, (f) as reagents for preparing probes in gene diagnoses, PCR primers, etc., (g) drugs for gene prophylactic and therapy, and the like.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION

(i) APPLICANT: Takeda Chemical Industries Ltd

(ii) TITLE OF THE INVENTION: Novel Human G-Protein Coupled Receptor Protein and its DNA

(iii) NUMBER OF SEQUENCES: 15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SmithKline Beecham, Corporate Intellectual Property

(B) STREET: Two New Horizons Court

(C) CITY: Brentford

(D) COUNTRY: Middlesex

(E) COUNTRY: United Kingdom

(F) POST CODE: TW8 9EP

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 286823/1996

(B) FILING DATE: 29 October 1996

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: CONNELL, Anthony Christopher

(B) GENERAL AUTHORISATION NUMBER:

(C) REFERENCE/DOCKET NUMBER: TAK50003

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: +44 127 964 4395

(B) TELEFAX: +44 181 975 6294

## (2) INFORMATION FOR SEQ ID NO: 1

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 481

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met Arg Trp Leu Trp Pro Leu Ala Val Ser Leu Ala Val Ile Leu Ala  
1 5 10 15Val Gly Leu Ser Arg Val Ser Gly Gly Ala Pro Leu His Leu Gly Arg  
20 25 30His Arg Ala Glu Thr Gln Gln Gln Ser Arg Ser Lys Arg Gly Thr  
35 40 45Glu Asp Glu Glu Ala Lys Gly Val Gln Gln Tyr Val Pro Glu Glu Trp  
50 55 60Ala Glu Tyr Pro Arg Pro Ile His Pro Ala Gly Leu Gln Pro Thr Lys  
65 70 75 80Pro Leu Val Ala Thr Ser Pro Asn Pro Asp Lys Asp Gly Gly Thr Pro  
85 90 95Asp Ser Gly Gln Glu Leu Arg Gly Asn Leu Thr Gly Ala Pro Gly Gln  
100 105 110Arg Leu Gln Ile Gln Asn Pro Leu Tyr Pro Val Thr Glu Ser Ser Tyr  
115 120 125Ser Ala Tyr Ala Ile Met Leu Leu Ala Leu Val Val Phe Ala Val Gly  
130 135 140Ile Val Gly Asn Leu Ser Val Met Cys Ile Val Trp His Ser Tyr Tyr  
145 150 155 160Leu Lys Ser Ala Trp Asn Ser Ile Leu Ala Ser Leu Ala Leu Trp Asp  
55

5	Phe Leu Val Leu Phe Cys Leu Pro Ile Val Ile Phe Asn Glu Ile	165	170	175	5	Asn Gln Phe Ser Thr Phe Phe Lys Gly Ala Ile Thr Pro Val Leu Leu	
10	Thr Lys Gln Arg Leu Leu Gly Asp Val Ser Cys Arg Ala Val Pro Phe	180	185	190	405	410	415
15	Met Glu Val Ser Ser Leu Gly Val Thr Thr Phe Ser Leu Cys Ala Leu	195	200	205	Leu Cys Ile Cys Arg Pro Leu Gly Gln Ala Phe Leu Asp Cys Cys Cys		
20	Gly Ile Asp Arg Phe His Val Ala Thr Ser Thr Leu Pro Lys Val Arg	210	215	220	420	425	430
25	Pro Ile Glu Arg Cys Gln Ser Ile Leu Ala Lys Leu Ala Val Ile Trp	225	230	235	240	Cys Cys Cys Cys Gln Gln Cys Gly Gly Ala Ser Glu Ala Ser Ala Ala	
30	Val Gly Ser Met Thr Leu Ala Val Pro Glu Leu Leu Trp Gln Leu	245	250	255	435	440	445
35	Ala Gln Glu Pro Ala Pro Thr Met Gly Thr Leu Asp Ser Cys Ile Met	260	265	270	450	455	460
40	Lys Pro Ser Ala Ser Leu Pro Glu Ser Leu Tyr Ser Leu Val Met Thr	275	280	285	Asn Gly Ser Asp Asn Lys Leu Lys Thr Glu Val Ser Ser Ser Ile Tyr		
45	Tyr Gln Asn Ala Arg Met Trp Trp Tyr Phe Gly Cys Tyr Phe Cys Leu	290	295	300	465	470	475
50	Pro Ile Leu Phe Thr Val Thr Cys Gln Leu Val Thr Trp Arg Val Arg	305	310	315	320	480	Cys
55	Gly Pro Pro Gly Arg Lys Ser Glu Cys Arg Ala Ser Lys His Glu Gln	325	330	335	(3) INFORMATION FOR SEQ ID NO: 2		
60	Ala Phe Cys Thr Leu Pro Glu Asn Val Cys Asn Ile Val Val Ala Tyr	340	345	350	(i) SEQUENCE CHARACTERISTICS:		
65	Leu Ser Thr Glu Leu Thr Arg Gln Thr Leu Asp Leu Leu Gly Leu Ile	355	360	365	(A) LENGTH: 542		
70		370	375	380	(B) TYPE: amino acid		
75		385	390	395	(C) STRANDEDNESS:		
80					(D) TOPOLOGY: linear		
85					(ii) MOLECULAR TYPE: peptide		
90					(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:		
95					Met Cys Pro Ala Glu Gly Pro Ala Arg Pro Val Ala Gly Gly Trp Glu		
100					1 5 10 15		
105					Gly Gly Gln Ala Ser Asp Ala Arg Arg Leu Thr Gly Gly Ser Ser		
110					20 25 30		
115					Arg Pro Ala Ala Ser Leu Glu Pro Ser Ser Trp Ala Pro Cys Thr His		
120					35 40 45		
125					Leu Leu Phe Leu Gly Trp Leu Ser Pro Ala His Pro Ala Met Arg Trp		
130					50 55 60		
135					Leu Trp Pro Leu Ala Val Ser Leu Ala Val Ile Leu Ala Val Gly Leu		
140					65 70 75 80		

Ser Arg Val Ser Gly Gly Ala Pro Leu His Leu Gly Arg His Arg Ala  
85 90 95

Glu Thr Gln Gln Gln Ser Arg Ser Lys Arg Gly Thr Glu Asp Glu  
100 105 110

Glu Ala Lys Gly Val Gln Gln Tyr Val Pro Glu Glu Trp Ala Glu Tyr  
115 120 125

Pro Arg Pro Ile His Pro Ala Gly Leu Gln Pro Thr Lys Pro Leu Val  
130 135 140

Ala Thr Ser Pro Asn Pro Asp Lys Asp Gly Gly Thr Pro Asp Ser Gly  
145 150 155 160  
Gln Glu Leu Arg Gly Asn Leu Thr Gly Ala Pro Gly Gln Arg Leu Gln  
165 170 175

Ile Gln Asn Pro Leu Tyr Pro Val Thr Glu Ser Ser Tyr Ser Ala Tyr  
180 185 190

Ala Ile Met Leu Leu Ala Leu Val Val Phe Ala Val Gly Ile Val Gly  
195 200 205

Asn Leu Ser Val Met Cys Ile Val Trp His Ser Tyr Tyr Leu Lys Ser  
210 215 220

Ala Trp Asn Ser Ile Leu Ala Ser Leu Ala Leu Trp Asp Phe Leu Val  
225 230 235 240  
Leu Phe Phe Cys Leu Pro Ile Val Ile Phe Asn Glu Ile Thr Lys Gln  
245 250 255

Arg Leu Leu Gly Asp Val Ser Cys Arg Ala Val Pro Phe Met Glu Val  
260 265 270

Ser Ser Leu Gly Val Thr Thr Phe Ser Leu Cys Ala Leu Gly Ile Asp  
275 280 285

Arg Phe His Val Ala Thr Ser Thr Leu Pro Lys Val Arg Pro Ile Glu  
290 295 300

Arg Cys Gln Ser Ile Leu Ala Lys Leu Ala Val Ile Trp Val Gly Ser

5

10

15

20

25

30

35

40

45

50

55

305 310 315 320  
Met Thr Leu Ala Val Pro Glu Leu Leu Trp Gln Leu Ala Gln Glu  
325 330 335

5

Pro Ala Pro Thr Met Gly Thr Leu Asp Ser Cys Ile Met Lys Pro Ser  
340 345 350

10

Ala Ser Leu Pro Glu Ser Leu Tyr Ser Leu Val Met Thr Tyr Gln Asn  
355 360 365

Ala Arg Met Trp Trp Tyr Phe Gly Cys Tyr Phe Cys Leu Pro Ile Leu  
370 375 380

15

Phe Thr Val Thr Cys Gln Leu Val Thr Trp Arg Val Arg Gly Pro Pro  
385 390 395 400  
Gly Arg Lys Ser Glu Cys Arg Ala Ser Lys His Glu Gln Cys Glu Ser  
405 410 415

20

Gln Leu Asn Ser Thr Val Val Gly Leu Thr Val Val Tyr Ala Phe Cys  
420 425 430

25

Thr Leu Pro Glu Asn Val Cys Asn Ile Val Val Ala Tyr Leu Ser Thr  
435 440 445

30

Glu Leu Thr Arg Gln Thr Leu Asp Leu Leu Gly Leu Ile Asn Gln Phe  
450 455 460

35

Ser Thr Phe Phe Lys Gly Ala Ile Thr Pro Val Leu Leu Cys Ile  
465 470 475 480  
Cys Arg Pro Leu Gly Gln Ala Phe Leu Asp Cys Cys Cys Cys Cys  
485 490 495

40

Cys Glu Glu Cys Gly Gly Ala Ser Glu Ala Ser Ala Asn Gly Ser  
500 505 510

45

Asp Asn Lys Leu Lys Thr Glu Val Ser Ser Ser Ile Tyr Phe His Lys  
515 520 525

50

Pro Arg Glu Ser Pro Pro Leu Leu Pro Leu Gly Thr Pro Cys  
530 535 540

55



## (4) INFORMATION FOR SEQ ID NO: 3

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1443

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

## (ii) MOLECULAR TYPE: cDNA

## (x) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

20 ATGGCGTGGC TGTGGCCCT GGCTGTCTCT CTTGCTGTGA TTTTGGCTGT  
 GGGGCTAAGC  
 60 AGGGTCTCTG GGGGTGCCCC CCTGCACCTG GGCAGGCACA GAGCCGAGAC  
 CCAGCGAGCAG  
 120 CAGAGCCGAT CCAAGAGGGG CACCGAGGAT GAGGAGGCCA  
 AGGGCGTGCA GCAGTATGT  
 180 CCTGAGGAGT GGGCGGAGTA CCCCCGGCC ATTCAACCTG CTGGCTGCA  
 GCCAACCAAG  
 240 CCCTTGGTGG CCACCAAGCC TAACCCCGAC AAGGATGGGG GCACCCAGA  
 CAGTGGGGCAG  
 300 GAACTGAGGG GCAATCTGAC AGGGGGACCA GGGCAGAGGC  
 TACAGATCCA GAACCCCTG  
 360 TATCCGGTGA CCGAGAGCTC CTACAGTGCC TATGCCATCA TCGTCTGGC  
 GCTGTGTGTG  
 420 TTTCGGGTGG GCA TTGTGGG CAACCTGTGG GTCATGTGCA TCGTGTGGCA  
 CAGCTACTAC  
 480 CTGAAGAGCG CCTGGAAC TC ATCTTTGCC AGCCTGGGCC TCTGGGATTT  
 TCTGTCTCTC  
 540 TTTTTCGCC TCCCTATTGT CATCTTCAAC GAGATCACC AAGCAGAGCT  
 ACTGGGTGAC  
 600

5 GTTTCCTGTC GTGCCGTGCC CTTCATGAG GTCTCTCTCTC TGGGAGTCAC  
 GACTTTCAGC  
 660 CTCTGTGCCC TGGGCATTGA CCGCTTCCAC GTGGCCACCA GCACCTGCC  
 CAAGGTGAGG  
 720 CCCATCGAGC GGTGCCAATC CATCTGGCC AAGTTGGCTG TCATCTGGGT  
 GGGCTCCATG  
 780 ACCGTGGCTG TGCTGAGCT CCTGTGTGG CAGCTGGCAG AGGAGCCTGC  
 CCCACCATG  
 840 GGCACCTGG ACTCATGCAI CATGAACCC TCAGCCAGCC TGCCCGAGTC  
 CCTGTATTCA  
 900 CTGTGATGA CCTACCAGAA CGCCCGCATG TGCTGTACT TTGGCTGCTA  
 CTCTGCGCTG  
 960 CCCATCTCT TCACAGTCAC CTGCCAGCTG GTGACATGGC GGGTGGAGG  
 CCTCCAGGG  
 1020 AGGAAGTCAG AGTGCAGGCG CAGCAAGCAG GAGCAGTGTG  
 AGAGCCAGCT CAACAGCACC  
 1080 GTGTGGGCC TGACCGTGTG CTACGCCCTTC TGACCCCTCC CAGAGAAGCT  
 CTGCAACATC  
 1140 GTGTGGCCT ACCTCTCCAC CGAGCTGACC CGCCAGACCC TGAACCTCT  
 GGGCTCATC  
 1200 AACCACTTCT CCACCTTCTT CAAAGGGCGCC ATCACCACCAG TGCTGTCTCT  
 TTGCATCTGC  
 1260 AGGCGCTGG GCCAGGCCCTT CTGGAGTGC TGCTGTCTGT GCTGTGTGA  
 GGAGTGGCGC  
 1320 GGGGCTTGG AGGCTCTGTC TGCCAAATGG TCGGACACA AGCTCAAGAC  
 CGAGGTGTC  
 1380 TCTTCATCT ACTTCACAA GCCCAGGGAG TCACCCCCAC TCCTGCCCC  
 GGGCACACCT

1440

TCC

1443

5

## (5) INFORMATION FOR SEQ ID NO: 4

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1626

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

10

## (ii) MOLECULAR TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

ATGTGTCAG CAGAGGGCCC TGCCGGCCCT GTGGCCGGAG GCTGGGAGGG  
AGGGCAGGCG

60

AGTGATGCCA GACGCCCTGAC TGGAGGCGGA TCCAGCCGGC CAGCTGCCTC  
TCTGGAGGCC

120

AGCTCTGGG CCCCCTGTAC TCACCTGCTC TTCCTGGGCT GGCTGTCTCC  
TGCTCATCCA

180

GCCATGCGGT GGCTGTGGCC CCTGGCTGTC TCCTTGCTG TGAITTTGGC  
TGTGGGGCTA

240

AGCAGGGTCT CTGGGGGTGC CCCCCTGCAC CTGGGCAGGC ACAGAGCCGA  
GACCCAGGAG

300

CAGCAGAGCC GATCCAAGAG GGGCACCGAG GATGAGGAGG  
CCAAAGGCGT GCAGCAGTAT

360

GTGCCCTGAGG AGTGGGCGGA GTACCCCGGG CCCATTCAAC CTGCTGGCCT  
GCAGCCCAACC

420

AAGCCCTTGG TGGCCACCAG CCCTAACCCC GACAAGGATG GGGGCAACCCC  
AGACAGTGGG

480

540 CAGGAACCTGA GGGGCAATCT GACAGGGGCA CCAGGGCAGA  
GGCTACAGAT CCAGAACCCC

55

CTGTATCCGG TGACCGAGAG CTCCTACAGT GCCTATGCCA TCATGCTTCT  
GGCGCTGGTG

600

GTGTTTGGG TGGCAATTGT GGGCAACCTG TCGGTCAATGT GCATCGTGTG  
GCACAGCTAC

660

TACCTGAAGA GCGCTGGAA CTCCATCCCTT GCCAGCCTGG CCCTCTGGGA  
TTTTCTGGTC

720

CTCTTTTCT GCCTCCCTAT TGTCACTTTC AACGAGATCA CCAAGCAGAG  
GCTACTGGGT

780

GACGTTTCT GTGCTGCCGT GCCCTTCAATG GAGGTCTCCT CTCTGGGAGT  
CACGACTTTC

840

AGCCTCTGTG CCCTGGGCAT TGACCGCTTC CACGTGGCCA CCAGCACCTT  
GCCCAAGGTG

900

AGGCCCATCG AGCGGTGCCA ATCCATCCTG GCCAAGTTGG CTGTCACTCTG  
GGTGGGCTCC

960

ATGACGCTGG CTGTGCCCTGA GCTCCTGCTG TGGCAGCTGG CACAGGAGCC  
TGCCCCCACC

1020

ATGGGCACCC TGGACTCATG CATCATGAAA CCCTCAGCCA GCCTGCCCGA  
GTCCCTGTAT

1080

TCACTGGTGA TGACCTACCA GAAAGCCCCG ATGTGGTGGT ACTTTGGCTG  
CTACTTCTGC

1140

CTGCCATCC TCTTCACAGT CACCTGCCAG CTGGTGACAT GGGGGGTGGG  
AGGCCCTCCA

1200

GGGAGGAAGT CAGAGTGCCAG GGCACGCAAG CACGAGCAGT  
GTGAGAGCCA GCTCAACAGC

1260

ACCGTGGTGG GCCTGACCGT GGTCTACGCC TTCTGCACCC TCCCAGAGAA  
CGTCTGCAAC

1320

ATCGTGTGG CCTACTCTC CACCGAGCTG ACCCGCCAGA CCTGGACCT  
CCTGGGCTC  
1380  
ATCAACCACT TCTCCACCTT CTTCAGGCG GCCATACCC CAGTGTCT  
CCTTGCATC  
1440  
TGCAGGCCCC TGGGCCAGGC CTTCCTGGAC TGCTGTGCT GCTGCTGTG  
TGAGGAGTGC  
1500  
GGCGGGCTT CGGAGGCTC TGCTGCCAAT GGGTCGACA ACAAGTCAA  
GACCGAGGTG  
1560  
TCCTCTTCCA TCTACTTCCA CAAGCCAGG GAGTACCCCC CACTCCTGCC  
CCTGGGCACA  
1620  
CCTTGC  
1626

(6) INFORMATION FOR SEQ ID NO: 5  
(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 426  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TGCAATCCAT CCTGGCCAAG TTGGGTGTA TCTGGGTGGG CTCCATGACG  
CTGGCTGTG  
60  
CTGAGCTCT GCTGTGGCAG CTGGCACAAG AGCCTGCCCC CACCATGGGG  
ACCTGGACT  
120  
CATGCATCAT GAAACCTCA GCCAGCCTGC CCGAGTCCCT GTATTCACTG  
GTGATGACCT  
180  
ACGAGAAGCG CCCCATGTGG TGGTACTTTG GCTGCTACTT CTGCTGCCCC  
ATCCTCTTA  
240

CAGTCACTGG CCAAGTGTG ACATGGCGGG TGCCAGGCCCC TCCAGGGAGG  
AAGTCAGAGT  
300  
GCAGGGCCAG CAAGCAGGAG CAGTGTGAGA GCCAGCTCAA CAGCACCTG  
GTGGGCTGA  
360  
CCGTGTCTA CGGCTTTTITG CAACCTTCA GAGAACGTTT GCAACATCGT  
GGTGGGCTTA  
420  
CCTTTT  
426

(7) INFORMATION FOR SEQ ID NO: 6  
(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 248  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AACAAGGCC GTGCTACG NCTTTCGAC CCTCCANAG AACGTCTGCA  
ACATCGTGT  
60  
GGCTTACCTC TCCACGAGC TGACCCGCA GNCCTGGAC CTCCTGGGCC  
TCATCAACCA  
120  
GTCTCAACC TTCTTCAAGG GCGCATCAC CCCAGTGCTG CTCCTTTGCA  
TCTGCAGGCC  
180  
GCTGGGCCAG GCTTCTCTG ACTGCTCTG CTGCTGCTGC TGTNAGGAGT  
GCGCGGGGC  
240  
TTGGGAGG  
248

(8) INFORMATION FOR SEQ ID NO: 7

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26

(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:  
AAGTTGGCTG TCATCTGGGT GGGCTC 26

(9) INFORMATION FOR SEQ ID NO: 8  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:  
TGAGCTCCTG CTGTGGCAGC TGGCAGC 28

(10) INFORMATION FOR SEQ ID NO: 9  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:  
CATGCGGGCG TTCTGGTAGG TCATCAC 27

(11) INFORMATION FOR SEQ ID NO: 10  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:  
GAAGAGGATG GGCAGGCAGA AGTAGCAG 28

(12) INFORMATION FOR SEQ ID NO: 11  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:  
ATGAAGGGCA CGGCAGCACA AGAAACG 27

(13) INFORMATION FOR SEQ ID NO: 12  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:  
ATGACAATAG GGAGGCAGAA AAAGAGG 27

(14) INFORMATION FOR SEQ ID NO: 13  
(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:  
GTCGACGAGA TGTGTGAGGG CAGCAAGAAG TGC 33

## SEQUENCE LISTING

5

(15) INFORMATION FOR SEQ ID NO: 14

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

TACTGGGGCC TCAGCAAGGT GTGCCACAG 28

20

(16) INFORMATION FOR SEQ ID NO: 15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GTCGACTGGC TGTCTCCTGC TCATCCAGCC AT 32

35

30

25

45

50

55

5

INFORMATION FOR SEQ ID NO: 1

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 481

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

20

Met Arg Trp Leu Trp Pro Leu Ala Val Ser Leu Ala Val Ile Leu Ala

1 5 10 15

Val Gly Leu Ser Arg Val Ser Gly Gly Ala Pro Leu His Leu Gly Arg

20 25 30

His Arg Ala Glu Thr Gln Glu Gln Ser Arg Ser Lys Arg Gly Thr

35 40 45

Glu Asp Glu Glu Ala Lys Gly Val Gln Gln Tyr Val Pro Glu Glu Trp

50 55 60

Ala Glu Tyr Pro Arg Pro Ile His Pro Ala Gly Leu Gln Pro Thr Lys

65 70 75 80

Pro Leu Val Ala Thr Ser Pro Asn Pro Asp Lys Asp Gly Gly Thr Pro

85 90 95

Asp Ser Gly Gln Glu Leu Arg Gly Asn Leu Thr Gly Ala Pro Gly Gln

100 105 110

55

5 Arg Leu Gln Ile Gln Asn Pro Leu Tyr Pro Val Thr Glu Ser Ser Tyr  
 115 120 125  
 Ser Ala Tyr Ala Ile Met Leu Leu Ala Leu Val Val Phe Ala Val Gly  
 130 135 140  
 10 Ile Val Gly Asn Leu Ser Val Met Cys Ile Val Thr Trp His Ser Tyr Tyr  
 145 150 155 160  
 15 Leu Lys Ser Ala Trp Asn Ser Ile Leu Ala Ser Leu Ala Leu Trp Asp  
 165 170 175  
 20 Phe Leu Val Leu Phe Phe Cys Leu Pro Ile Val Ile Phe Asn Glu Ile  
 180 185 180  
 Thr Lys Gln Arg Leu Leu Gly Asp Val Ser Cys Arg Ala Val Pro Phe  
 195 200 205  
 25 Met Glu Val Ser Ser Leu Gly Val Thr Thr Phe Ser Leu Cys Ala Leu  
 210 215 220  
 30 Gly Ile Asp Arg Phe His Val Ala Thr Ser Thr Leu Pro Lys Val Arg  
 225 230 235 240  
 35 Pro Ile Glu Arg Cys Gln Ser Ile Leu Ala Lys Leu Ala Val Ile Trp  
 245 250 255  
 40 Val Gly Ser Met Thr Leu Ala Val Pro Glu Leu Leu Leu Trp Gln Leu  
 260 265 270  
 45 Ala Gln Glu Pro Ala Pro Thr Met Gly Thr Leu Asp Ser Cys Ile Met  
 275 280 285  
 50  
 55

5 Lys Pro Ser Ala Ser Leu Pro Glu Ser Leu Tyr Ser Leu Val Met Thr  
 290 295 300  
 Tyr Gln Asn Ala Arg Met Trp Trp Tyr Phe Gly Cys Tyr Phe Cys Leu  
 305 310 315 320  
 10 Pro Ile Leu Phe Thr Val Thr Cys Gln Leu Val Thr Trp Arg Val Arg  
 325 330 335  
 15 Gly Pro Pro Gly Arg Lys Ser Glu Cys Arg Ala Ser Lys His Glu Gln  
 340 345 350  
 20 Cys Glu Ser Gln Leu Asn Ser Thr Val Val Gly Leu Thr Val Val Tyr  
 355 360 365  
 Ala Phe Cys Thr Leu Pro Glu Asn Val Cys Asn Ile Val Val Ala Tyr  
 370 375 380  
 25 Leu Ser Thr Glu Leu Thr Arg Gln Thr Leu Asp Leu Leu Gly Leu Ile  
 385 390 395 400  
 30 Asn Gln Phe Ser Thr Phe Phe Lys Gly Ala Ile Thr Pro Val Leu Leu  
 405 410 415  
 35 Leu Cys Ile Cys Arg Pro Leu Glu Gln Ala Phe Leu Asp Cys Cys  
 420 425 430  
 40 Cys Cys Cys Cys Glu Glu Cys Gly Ala Ser Glu Ala Ser Ala Ala  
 435 440 445  
 45 Asn Gly Ser Asp Asn Lys Leu Lys Thr Glu Val Ser Ser Ser Ile Tyr  
 450 455 460  
 Phe His Lys Pro Arg Glu Ser Pro Pro Leu Leu Pro Leu Gly Thr Pro  
 465 470 475 480  
 50 Cys  
 55

## INFORMATION FOR SEQ ID NO: 2

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 542  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (11) MOLECULAR TYPE: peptide

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Cys Pro Ala Glu Gly Pro Ala Arg Pro Val Ala Gly Gly Trp Glu  
 1 5 10 15  
 Gly Gly Gln Ala Ser Asp Ala Arg Arg Leu Thr Gly Gly Ser Ser  
 20 25 30  
 Arg Pro Ala Ala Ser Leu Glu Pro Ser Ser Trp Ala Pro Cys Thr His  
 35 40 45  
 Leu Leu Phe Leu Gly Trp Leu Ser Pro Ala His Pro Ala Met Arg Trp  
 50 55 60  
 Leu Trp Pro Leu Ala Val Ser Leu Ala Val Ile Leu Ala Val Gly Leu  
 65 70 75 80  
 Ser Arg Val Ser Gly Gly Ala Pro Leu His Leu Gly Arg His Arg Ala  
 85 90 95  
 Glu Thr Gln Glu Gln Ser Arg Ser Lys Arg Gly Thr Glu Asp Glu  
 100 105 110  
 Glu Ala Lys Gly Val Gln Gln Tyr Val Pro Glu Glu Trp Ala Glu Tyr  
 115 120 125  
 Pro Arg Pro Ile His Pro Ala Gly Leu Gln Pro Thr Lys Pro Leu Val  
 130 135 140  
 Ala Thr Ser Pro Asn Pro Asp Lys Asp Gly Gly Thr Pro Asp Ser Gly  
 145 150 155 160

Gln Glu Leu Arg Gly Asn Leu Thr Thr Gly Ala Pro Gly Gln Arg Leu Gln  
 165 170 175  
 Ile Gln Asn Pro Leu Tyr Pro Val Thr Glu Ser Ser Tyr Ser Ala Tyr  
 180 185 190  
 Ala Ile Met Leu Leu Ala Leu Val Val Phe Ala Val Gly Ile Val Gly  
 195 200 205  
 Asn Leu Ser Val Met Cys Ile Val Trp His Ser Tyr Tyr Leu Lys Ser  
 210 215 220  
 Ala Trp Asn Ser Ile Leu Ala Ser Leu Ala Leu Trp Asp Phe Leu Val  
 225 230 235 240  
 Leu Phe Phe Cys Leu Pro Ile Val Ile Phe Asn Glu Ile Thr Lys Gln  
 245 250 255  
 Arg Leu Leu Gly Asp Val Ser Cys Arg Ala Val Pro Phe Met Glu Val  
 260 265 270  
 Ser Ser Leu Gly Val Thr Thr Phe Ser Leu Cys Ala Leu Gly Ile Asp  
 275 280 285  
 Arg Phe His Val Ala Thr Ser Thr Leu Pro Lys Val Arg Pro Ile Glu  
 290 295 300  
 Arg Cys Gln Ser Ile Leu Ala Lys Leu Ala Val Ile Trp Val Gly Ser  
 305 310 315 320  
 Met Thr Leu Ala Val Pro Glu Leu Leu Trp Gln Leu Ala Gln Glu  
 325 330 335  
 Pro Ala Pro Thr Met Gly Thr Leu Asp Ser Cys Ile Met Lys Pro Ser  
 340 345 350  
 Ala Ser Leu Pro Glu Ser Leu Tyr Ser Leu Val Met Thr Tyr Gln Asn  
 355 360 365

Ala Arg Met Trp Trp Trp Phe Gly Cys Tyr Phe Cys Leu Pro Ile Leu  
 370 375 380  
 Phe Thr Val Thr Cys Gln Leu Val Thr Trp Arg Val Arg Gly Pro Pro  
 385 390 395 400  
 Gly Arg Lys Ser Glu Cys Arg Ala Ser Lys His Glu Gln Cys Glu Ser  
 405 410 415  
 Gln Leu Asn Ser Thr Val Val Gly Leu Thr Val Val Tyr Ala Phe Cys  
 420 425 430  
 Thr Leu Pro Glu Asn Val Cys Asn Ile Val Val Ala Tyr Leu Ser Thr  
 435 440 445  
 Glu Leu Thr Arg Gln Thr Leu Asp Leu Leu Gly Leu Ile Asn Gln Phe  
 450 455 460  
 Ser Thr Phe Phe Lys Gly Ala Ile Thr Pro Val Leu Leu Cys Ile  
 465 470 475 480  
 Cys Arg Pro Leu Gly Gln Ala Phe Leu Asp Cys Cys Cys Cys Cys  
 485 490 495  
 Cys Glu Glu Cys Gly Gly Ala Ser Glu Ala Ser Ala Ala Asn Gly Ser  
 500 505 510  
 Asp Asn Lys Leu Lys Thr Glu Val Ser Ser Ser Ile Tyr Phe His Lys  
 515 520 525  
 Pro Arg Glu Ser Pro Pro Leu Leu Pro Leu Gly Thr Pro Cys  
 530 535 540

## INFORMATION FOR SEQ ID NO: 3

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1443  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (ii) MOLECULAR TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATGCGGTGGC TGTGGCCCT GCGTGTCTCT CTGCTGTGCA TTTGGCTGT GGGGCTAAGC 60  
 AGGTCTCTG GGGTGGCCC CCTGCACCTG GGCAGGCACA GAGCGAGAC CCAGGAGCAG 120  
 CAGAGCCGAT CCAAGAGGGG CACCGAGGAT GAGGAGGCCA AGGGCTGCA GCAGTATGTG 180  
 CCTGAGGACT GGGCGGAGTA CCCCCGGCC ATTACCCCTG CTGGCTCCA GCCAACCAAG 240  
 CCCTTGTGG CCACCAAGCC TAACCCGAC AAGGATGGG GCACCCAGA CAGTGGCAG 300  
 GAACTGAGG GCAATCTGAC AGGGCACCA GGGCAGGC TACAGATCCA GAACCCCTG 360  
 TATCCGGTGA CCGAGAGCTC CTACAGTGGC TATGCCATCA TGGTTGGG GGTGTGTG 420  
 TTTCGGTGG GCATTGGG CAACCTGTG GTCATGTGCA TCGTGTGCA CAGCTACTAC 480



CTGAAAGGG CCTGAACTC CATCTTCC ACCCTGGCC CTGGGAATT TCTGTCCTC 540  
TTTTTGGC TCCCAATGT CATCTTAC GAGATACCA AGCAAGGCT ACTGGGTAC 600  
GTTCCTGTG GTCCCTGCC CTTCATGAG GTCTCTCTC TGGAGTAC GACTTTACG 660  
CTCTGCCC TGGCAATTA CCGCTTCC AC GTGGCACA GCACCTGCC CAAGTGAGG 720  
CCCATGAGG GTTCCCAATC CATCTGGCC AAGTTGGTG TCAATGGGT GGGCTGCAATG 780  
ACCTGGCTG TCCCTAGCT CTGTGTGTG CAGCTGGCA AGCAAGCTGC CCCACCAATG 840  
GGCAACCTGG ACTATGCAI CATGAACCC TACGCCAGC TCCCGAGTC CCTGAATTA 900  
CTGTGATGA CCTACAGAA GCGCCGCAATG TGTGTACT TTGGCTGCTA CTCTGCTG 960  
CCCATCTCT TCAAGTAC CTGCAAGCT GTGACATGG GGTGGGAGG CCTTCAGGG 1020  
AGCAAGTCA AGTCAAGGG CAGAAAGCA GAGCACTGT AGACCAAGT CAACAGCAC 1080  
GTGTGGGG TGAAGTGT CTACGCTTC TCAAGCTGC CAGAAAGCT CTGCAATC 1140  
GTGTGGCT ACTCTGCA CAGCTGACC GGCAGAGCC TGAAGTCT GGGCTCATC 1200  
AACAGTCT CAGCTTCT CAAGGGGGC ATACCCAG TGTGTCTCT TTGCAATG 1260  
AGCCGCTGG CCAAGGCTT CTGTAGTGC TGTGTCTCT CCTCTGTGA GGAATGCGG 1320  
GGGCTTGG AGGCTCTGC TGCATGGG TCGAACAACA AGCTCAAGAC CAAAGTGTG 1380  
TCTTCAATC ACTTCAACA CCGCAGGAG TCAACCCCA TCTGCCCCT GGCACACCT 1440  
TGC 1443

## INFORMATION FOR SEQ ID NO: 4

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1626  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: cDNA

(12) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

ATGTGCAAG CAGAGGGCC TCCCGGCT GTGGCCGAG CCTGGAGCG AGGCAGCG 60  
AGTATGCCA GACCCCTGAC TGAAGCGGA TCCAGCCGG CAGCTGGCT TCTGAACCC 120  
AGCTTGGG CCCCCTGAC TCACTGCTC TTCTGGGT GGTGTCTCC TCTCATCA 180  
GCCATGCGT GGTGTGGC CTTGGCTGC TCTTCTCT TATTTGGC TGTGGGCTA 240

## INFORMATION FOR SEQ ID NO: 5

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 426  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

## (ii) MOLECULAR TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

5 ACCAGGCTCT CTGGGGTGG CCCCTGGAC CTGGGCGAGG ACAGGCCCA GACCCAGGAG 300  
CAGCAGAGCC GATCCAGAG GGGCAGCGAG GATGAGGAG CCAAGGGCTT GCAGCAGTAT 360  
GTGGCTCAGG AGTGGGCGA GTACCCCGGG GCCATTACCC CTGTGGCTCT GCAGGCCAAC 420  
AAGCCCTTGG TGGCCACCAAG CCCATACCCC GACAAGGATG GGGGCACCCC AGACAGTGGG 480  
CAGGAACCTGA GGGGCAATCT CACAGGGGCA CCAGGGCAGA GGCTACAGAT CCAGAACCCC 540  
CTGTATCCGG TGACGAGAG CTCTATAGAT GCGTATGCCA TCAATCTTCT GGGCTGGTG 600  
GTGTTTGGGG TGGGCAATCT GGGCAACCTG TCGGTCACTT GCAATGGTGG GCACAGCTAC 660  
TACCTGAAGA GGGGCTGGA CTCATCTCTT GCCAGGCTGG CCGTCTGGGA TTTTCTGGTC 720  
CTCTTTTCTT GCGTCCCTAT TGTATCTTC AACGAGATCA CCAAGCAGAG GGTACTGGGT 780  
GAGCTTTCTT CTGTGGCGGT GCGCTTCATG GAGTCTCTCT CTCTGGGAGT CACGACTTTC 840  
AGGCTCTGTG CCGTGGGCAI TGACCGGCTC CAGTGGGCA CCAGCACCTT GCGCAAGGTG 900  
AGGCCCATGG AGGGGTGCCA ATGCATCCCTG GCGAAGTTGG CTGTCACTGG GGTGGGCTCC 960  
ATGACCTGG CTGTGGCTGA GCTCTCTGTG TGGCAGCTGG CACAGAGGCC TGGCGGCACC 1020  
ATGGGCACCC TGGACTCATG CATCATGAA CCGTCAGCCA GCGTGGCGGA GTCCCTGTAT 1080  
TCACTGGTGA TGACCTACCA GAAGCGCCGC ATGTGGTGGT ACTTGGCTG CTACTTCTCC 1140  
CTGCCCATCC TCTTACAGT CAGCTGGCAG CTGGTGACAT GGGGGGTGG AGCGCTCCA 1200  
GGGAGGAAGT CAGAGTCCAG GGGCAGCAG CAGGAGCAGT GTGAGAGCCA GGTCAACAGC 1260  
ACCGTGGTGG CCGTAGCGGT GGTCTAGCC TTCTGACCC TCCACAGAA GGTCTGGCAC 1320  
ATGGTGGTGG CCGTACTCTC CACGAGCTG ACCGGCCAGA CCGTGGACCT CTTGGGCTC 1380  
ATCAACCAAT TCTCCACCTT CTCAAGGGC GCGATCACCC CAGTGTCTGT CTTTGGCATC 1440  
TGCAGGGCGG TGGGCCAGGC CTTCCTGGAC TCGTGTGGT GCTGTGGTGT TGAGGAGTGG 1500  
GGCGGGGCTT CCGAGGCGTC TCGTGGCAAT GGGTGGGACA ACAAGCTCAA GACCGAGGTG 1560  
TCCGTTTCCA TCTACTCCA CAAGGCGAGG GAGTCAACCC CACTCTGGCC CTTGGGCACA 1620  
CGTTGC 1628

## INFORMATION FOR SEQ ID NO: 6

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 248  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: cDNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

ACGAAGGGCC GTGCTCTACG NCTTCTGCAC CGTCCGACAG AACCTCTCCA ACATCTGTGT 80  
GGCCTACCTC TCCAGCGAGC TGACCCGGCA GNCCTTGAC CTCTGGGCC TCATCAACGA 120  
GTCTCCACC TTCTTCAGG GCGGCATGAC CCAAGTGCTG CTCTTTGCA TGTCAAGCCC 180  
GCTGGGCCAG GCTTCTCTGG ACTGCTGCTG CTGCTGCTGC TGTAGAGAGT CCGCGCGGGC 240  
TTCCGAGG 248

## INFORMATION FOR SEQ ID NO: 7

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: synthetic DNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

AAGTTGGCTG TCATCTGGCT GGCTTC 26

## INFORMATION FOR SEQ ID NO: 8

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: synthetic DNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TGAGTCTCTG CTGTGGCAGC TGCACAG 28

## INFORMATION FOR SEQ ID NO: 9

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(11) MOLECULAR TYPE: synthetic DNA

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CATGGGGCGG TTCTGTAGG TCATCAC 27

INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GAAGAGGATG GGCAGGCAGA AGTAGCAG 28

INFORMATION FOR SEQ ID NO: 11

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ATGAGGGCA GGCAGGCAGA AGAAGC 27

INFORMATION FOR SEQ ID NO: 12

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

ATGACAATAG GGAGGCAGAA AAGAGC 27

INFORMATION FOR SEQ ID NO: 13

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GTCGACGAGA TGTGTGAGGG CAGCAAGAG TGC 33

INFORMATION FOR SEQ ID NO: 14

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

TACTGGGGC TCAGCAAGGT GTGCCAG 28

INFORMATION FOR SEQ ID NO: 15

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 32  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULAR TYPE: synthetic DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GTCCAGCTGGC TGTCTCTGC TCATCCAGCC AT 32

## Claims

1. A G-protein coupled receptor protein which comprises the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 1, or a variant of the amino acid sequence having a deletion, addition or

substitution of 1 to 30 amino acids, or its salt.

2. The G-protein coupled receptor protein of claim 1 which comprises the same or substantially the same amino acid sequence as that represented by SEQ ID NO: 2, or a variant of the amino acid sequence having a deletion, addition or substitution of 1 to 30 amino acids, or its salt.

3. A partial peptide of the G-protein coupled receptor protein of claim 1 or its salt.

4. An isolated DNA comprising DNA having a nucleotide sequence encoding the G-protein coupled receptor protein of claim 1.

5. The isolated DNA of claim 4 having the nucleotide sequence represented by SEQ ID NO: 3.

6. The isolated DNA of claim 4 having the nucleotide sequence represented by SEQ ID NO: 4.

A recombinant vector comprising the DNA of claim 4.

8. A transformant comprising the recombinant vector of claim 7.

9. The transformant of claim 8 which is *Escherichia coli* HB101/pHEBR2 (FERM BP-5724).

10. A process for preparing the G-protein coupled receptor protein of claim 1 or its salt which comprises cultivating the transformant of claim 8 to form the G-protein coupled receptor protein and recovering it.

11. A method for determining a ligand to the G-protein coupled receptor protein of claim 1 or its salt which comprises bringing the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt into contact with a test compound.

12. A method for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of claim 1 or its salt, or their salts which comprises comparing

(i) ligand binding upon bringing the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt into contact with the ligand, and  
(ii) that upon bringing the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt into contact with the ligand and a test compound.

13. The screening method according to claim 12, wherein ligand binding is compared by means of an amount of a labeled ligand bound to the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt; an amount of a labeled ligand bound to cells containing the G-coupled receptor protein or a membrane fraction of cells containing the G-protein coupled receptor protein of claim 1; an amount of a labeled ligand bound to the G-protein coupled receptor protein of claim 1 expressed on a cell membrane of the transformant of claim 8; or cell stimulation activities mediated by the G-protein coupled receptor protein of claim 1.

14. A kit for screening for compounds which alter binding of a ligand to the G-protein coupled receptor protein of claim 1 or its salt, or their salts which comprises as an essential component the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt.

15. The kit of claim 14, wherein the component is in the form of cells containing the G-protein coupled receptor protein of claim 1 or its salt or a cell membrane fraction of cells containing the G-protein coupled receptor protein of claim 1 or its salt.

16. A compound which alters ligand binding to the G-protein coupled receptor protein of claim 1 or its salt obtained by the screening method of claim 12 or the kit of claim 14.

17. An antibody against the G-protein coupled receptor protein of claim 1 or its salt or the partial peptide of claim 3 or its salt.

18. A method for determining the G-protein coupled receptor protein of claim 1, or its salt or the partial peptide of claim

3 or its salt in a specimen which comprises bringing the antibody of claim 17 into contact with the G-protein coupled receptor protein of claim 1, the partial peptide of claim 3 or a salt thereof in the specimen.

19. A pharmaceutical composition comprising as an effective component the G-protein coupled receptor protein of claim 1, or its salt or the partial peptide of claim 3 or its salt, and a pharmaceutically acceptable carrier or diluent.

20. A pharmaceutical composition comprising as an effective component the compound of claim 16 or its salt, and a pharmaceutically acceptable carrier or diluent.











Fig. 5

